

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**

## Yeast microarrays for genome wide parallel genetic and gene expression analysis

DEVAL A. LASHKARI\*<sup>†</sup>, JOSEPH L. DERISI<sup>‡</sup>, JOHN H. MCCUSKER<sup>§</sup>, ALLEN F. NAMATH<sup>‡</sup>, CRISTL GENTILE<sup>§</sup>, SEUNG Y. HWANG<sup>‡</sup>, PATRICK O. BROWN<sup>‡</sup>, AND RONALD W. DAVIS\*<sup>‡</sup>

Departments of \*Genetics and <sup>‡</sup>Biochemistry, Stanford University, Stanford, CA 94305; and <sup>§</sup>Department of Microbiology, Duke University, Durham, NC 27710

Contributed by Ronald W. Davis, September 2, 1997

**ABSTRACT** We have developed high-density DNA microarrays of yeast ORFs. These microarrays can monitor hybridization to ORFs for applications such as quantitative differential gene expression analysis and screening for sequence polymorphisms. Automated scripts retrieved sequence information from public databases to locate predicted ORFs and select appropriate primers for amplification. The primers were used to amplify yeast ORFs in 96-well plates, and the resulting products were arrayed using an automated microarraying device. Arrays containing up to 2,479 yeast ORFs were printed on a single slide. The hybridization of fluorescently labeled samples to the array were detected and quantitated with a laser confocal scanning microscope. Applications of the microarrays are shown for genetic and gene expression analysis at the whole genome level.

The genome sequencing projects have generated and will continue to generate enormous amounts of sequence data. The genomes of *Saccharomyces cerevisiae*, *Haemophilus influenzae* (1), *Mycoplasma genitalium* (2), and *Methanococcus jannischii* (3) have been completely sequenced. Other model organisms have had substantial portions of their genomes sequenced as well including the nematode *Caenorhabditis elegans* (4) and the small flowering plant *Arabidopsis thaliana* (5). Given this ever-increasing amount of sequence information, new strategies are necessary to efficiently pursue the next phase of the genome projects—the elucidation of gene expression patterns and gene product function on a whole genome scale.

One important use of genome sequence data is to attempt to identify the functions of predicted ORFs within the genome. Many of the ORFs identified in the yeast genome sequence were not identified in decades of genetic studies and have no significant homology to previously identified sequences in the database. In addition, even in cases where ORFs have significant homology to sequences in the database, or have known sequence motifs (e.g., protein kinase), this is not sufficient to determine the actual biological role of the gene product. Experimental analysis must be performed to thoroughly understand the biological function of a given ORF's product. Model organisms, such as *S. cerevisiae*, will be extremely important in improving our understanding of other more complex and less manipulable organisms.

To examine in detail the functional role of individual ORFs and relationships between genes at the expression level, this work describes the use of genome sequence information to study large numbers of genes efficiently and systematically. The procedure was as follows. (i) Software scripts scanned annotated sequence information from public databases for predicted ORFs. (ii) The start and stop position of each identified ORF was extracted automatically, along with the sequence data of the ORF and 200

bases flanking either side. (iii) These data were used to automatically select PCR primers that would amplify the ORF. (iv) The primer sequences were automatically input into the automated multiplex oligonucleotide synthesizer (6). (v) The oligonucleotides were synthesized in 96-well format, and (vi) used in 96-well format to amplify the desired ORFs from a genomic DNA template. (vii) The products were arrayed using a high-density DNA arrayer (7–10). The gene arrays can be used for hybridization with a variety of labeled products such as cDNA for gene expression analysis or genomic DNA for strain comparisons, and genomic mismatch scanning purified DNA for genotyping (11).

### METHODS

**Script Design.** All scripts were written in UNIX Tool Command Language. Annotated sequence information from GenBank was extracted into one file containing the complete nucleotide sequence of a single chromosome. A second file contained the assigned ORF name followed by the start and stop positions of that ORF. The actual sequence contained within the specified range, along with 200 bases of sequence flanking both sides, was extracted and input into the primer selection program PRIMER 0.5 (Whitehead Institute, Boston). Primers were designed so as to allow amplification of entire ORFs. The selected primer sequences were read by the 96-well automated multiplex oligonucleotide synthesizer instrument for primer synthesis. The forward and reverse primers were synthesized in two separate 96-well plates in corresponding wells. All primers were synthesized on a 20-nmol scale.

**ORF Amplification and Purification.** Genomic DNA was isolated as described (12) and used as template for the amplification reactions. Each PCR was done in a total volume of 100  $\mu$ l. A total of 0.2  $\mu$ M each of forward and reverse primers were aliquoted into a 96-well PCR plate (Robbins Scientific, Sunnyvale, CA); a master mix containing 0.24 mM each dNTP, 10 mM Tris (pH 8.5), 50 mM MgCl<sub>2</sub>, 2.5 units *Taq* polymerase, and 10 ng of template was added to the primers, and the entire mix was thermal cycled for 30 cycles as follows: 15 min at 94°C, 15 min at 54°C, and 30 min at 72°C. Products were ethanol precipitated in polystyrene v-bottom 96-well plates (Costar). All samples were dried and stored at –20°C.

**Arraying Procedure and Processing.** Microarrays were made as described (8).

A custom built arraying robot was used to print batches of 48 slides. The robot utilizes four printing tips which simultaneously pick up  $\approx$ 1  $\mu$ l of solution from 96-well microtiter plates. After printing, the microarrays were rehydrated for 30 sec in a humid chamber and then snap dried for 2 sec on a hot plate (100°C). The DNA was then UV crosslinked to the surface by subjecting the slides to 60 millijoules of energy. The rest of the poly-L-lysine surface was blocked by a 15-min incubation in a solution of 70 mM succinic anhydride dissolved in a solution consisting of 315 ml of 1-methyl-2-pyrrolidinone (Aldrich) and 35 ml of 1 M boric acid (pH 8.0). Directly after the blocking reaction, the bound DNA was denatured by a 2-min incubation in distilled water at  $\approx$ 95°C.

The publication costs of this article were defrayed in part by page charge payment. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. §1734 solely to indicate this fact.

© 1997 by The National Academy of Sciences 0027-8424/97/9413057-6\$2.00/0  
PNAS is available online at <http://www.pnas.org>.

Abbreviation: YEP, yeast extract/peptone.

<sup>†</sup>To whom reprint requests should be sent at the present address: Synteni, Inc., 6519 Dumbarton Circle, Fremont, CA 94555.




FIG. 1. Two-color fluorescent scan of a yeast microarray containing 2,479 elements (ORFs). The center-to-center distance between elements is 345  $\mu\text{m}$ . A probe mixture consisting of cDNA from yeast extract/peptone (YEP) galactose (green pseudocolor) and YEP glucose (red pseudocolor) grown yeast cultures was hybridized to the array. Intensity per element corresponds to ORF expression, and pseudocolor per element corresponds to relative ORF expression between the two cultures.

The slides were then transferred into a bath of 100% ethanol at room temperature.

**Probe Preparation: cDNA.** Yeast cultures (100 ml) were grown to  $\sim 1$  OD<sub>600</sub> and total RNA was isolated as described (13). Up to 500  $\mu\text{g}$  total RNA was used to isolate mRNA (Qiagen, Chatsworth, CA). Oligo(dT)20 (5  $\mu\text{g}$ ) was added and annealed to 2  $\mu\text{g}$  of mRNA by heating the reaction to 70°C for 10 min and quick chilling on ice, plus 2  $\mu\text{l}$  SuperScript II (200 units/ $\mu\text{l}$ ) (Life Technologies, Gaithersburg, MD), 0.6  $\mu\text{l}$  50 $\times$  dNTP mix (final concentrations were 500  $\mu\text{M}$  dATP, dCTP, dGTP, and 200  $\mu\text{M}$  dTTP), 6  $\mu\text{l}$  5 $\times$  reaction buffer, and 60  $\mu\text{M}$  Cy3-dUTP or Cy5-dUTP (Amersham). Reactions were carried out at 42°C for 2 h, after which the mRNA was degraded by the addition of 0.3  $\mu\text{l}$  5 M NaOH and 0.3  $\mu\text{l}$  100 mM EDTA and heating to 65°C for 10 min. The sample was then diluted to 500  $\mu\text{l}$  with TE and concentrated using a Microcon-30 (Amicon) to 10  $\mu\text{l}$ .

**Probe Preparation: Genomic DNA.** Fluorescent DNA was prepared from total genomic DNA as follows: 1  $\mu\text{g}$  of random nonamer oligonucleotides was added to 2.5  $\mu\text{g}$  of genomic DNA. This mixture was boiled for 2 min and then chilled on ice. A reaction mixture containing dNTPs (25  $\mu\text{M}$  dATP, dCTP, dGTP, 10  $\mu\text{M}$  dTTP, and 40  $\mu\text{M}$  Cy3-dUTP or Cy5-dUTP) reaction buffer (New England Biolabs), and 20 units exonuclease free Klenow enzyme (United States Biochemical) was added, and the reaction was incubated at 37°C for 2 h. The sample was then diluted to 500  $\mu\text{l}$  with TE and concentrated using a Microcon-30 (Amicon) to 10  $\mu\text{l}$ .

**Hybridization.** Purified, labeled probe was resuspended in 11  $\mu\text{l}$  of 3.5 $\times$  SSC containing 10  $\mu\text{g}$  *Escherichia coli* tRNA, and 0.3% SDS. The sample was then heated for 2 min in boiling water, cooled rapidly to room temperature, and applied to the array. The array was placed in a sealed, humidified, hybridization chamber. Hybridization was carried out for 10 h in a 62°C water bath, after which the arrays were washed immediately in 2 $\times$  SSC/0.2% SDS. A second wash was performed in 0.1 $\times$  SSC.

**Analysis and Quantitation.** Arrays were scanned on a scanning laser fluorescence microscope developed by Steve Smith with software written by Noam Ziv (Stanford Univer-

sity). A separate scan was done for each of the two fluorophores used. The images were then combined for analysis. A bounding box, fitted to the size of the DNA spots, was placed over each array element. The average fluorescent intensity was calculated by summing the intensities of each pixel present in a bounding box and then dividing by the total number of pixels. Local area background was calculated for each array element by determining the average fluorescent intensity at the edge of the bounding box. To normalize for fluorophore-specific variation, control spots containing yeast genomic DNA were applied to each quadrant during the arraying process. These elements were quantitated and the ratios of the signals were determined. These ratios were then used to normalize the photomultiplier sensitivity settings such that the ratios of the fluorescence of the genomic DNA spots were close to a value of 1.0. The average signal intensity at any given spot was regarded as significant if it was at least two standard deviations above background. Each experiment was conducted in duplicate, with the fluorophores representing each channel reversed. The ratios presented here are the average of the two experiments, except in the case in which the signal for the element in question was below the reliability threshold. The reliability threshold also determined the dynamic range of the experiment. For all of the experiments presented, the average dynamic range was  $\sim 1$  to 100. In the case where the fluorescence from a very bright spot saturates the detector, differential ratios will, in general, be underestimated. This can be compensated for by scanning at a lower overall sensitivity.

## RESULTS

The accumulation of sequence information from model organisms presents an enormous opportunity and challenge to understand the biological function of many previously uncharacterized genes. To do this accurately and efficiently, a directed strategy was developed that enables the monitoring of multiple genes simultaneously. Microarraying technology provides a method by which DNA can be attached to a glass surface in a high-density format (8). In practice, it is possible to array over 6,000 elements in an area less than 1.8 cm<sup>2</sup>. Given that the yeast genome consists of  $\sim 6,100$  ORFs, the entire set of yeast genes can be spotted onto a single glass slide.

With this capability and the availability of the entire sequence of the yeast genome, our strategy was to use a directed approach for generating the complete genome array. This procedure involved synthesizing a pair of oligonucleotide primers to amplify each ORF. The PCR product containing each gene of interest was arrayed onto glass and used, for example, as probe for monitoring gene expression levels by hybridizing to the array labeled cDNA generated from isolated mRNA of a culture grown under any experimental condition.

**Primer Selection and Synthesis.** The primer selection was fully automated using Tool Command Language scripts and PRIMER 0.5. (Whitehead). Primer pairs were automatically selected successfully for >99% of the ORFs tested. Primer sequences can thus be selected rapidly with minimal manual processing. A complete set of forward and reverse primers were selected initially for each ORF on chromosomes I, II, III, V, VI, VIII, IX, X, and XI. Primers for a representative set of ORFs (15% coverage) were chosen for the remaining chromosomes. With the release of the entire yeast genome sequence, the complete set of primers has now been selected.

Because each ORF requires a unique pair of synthetic primers, a total of approximately 12,200 oligonucleotides will be required to individually amplify each target. This costly component was addressed with the automated multiplex oligonucleotide synthesizer (6) which efficiently synthesizes primers in a 96-well format. Each primer, synthesized on a 20-nmol scale, provides enough material for 100 amplification reactions, whereas a given PCR product provides enough material to generate an element on

Table 1. Heat shock vs. control expression data

Ratio of gene expression		ORF	Gene	Description
Control	Heat			
2.3	2.2	YLR142	PUT1	Proline oxidase
	2.0	YOL140	ARG8	Acetylornithine aminotransferase
		YGL148	ARO2	Chorismate synthase
	36.0	YFL014	HSP12	Heat shock protein
	27.4	YBR072	HSP26	Heat shock protein
	6.7	YBR054	YRO2	Similarity to HSP30 heat shock protein Yrolp
	3.4	YCR021	HSP30	Heat shock protein
	2.6	YER103	SSA4	Heat shock protein
	2.5	YLR259	HSP60	Mitochondrial heat shock protein HSP60
	2.1	YBR169	SSE2	Heat shock protein of the HSP70 family
	1.7	YBL075	SSA3	Cytoplasmic heat shock protein
	1.4	YPL240	HSP82	Heat shock protein
	1.4	YDR258	HSP78	Mitochondrial heat shock protein of clpb family of ATP-dependent proteases
1.0		YNL007	SIS1	Heat shock protein
1.1		YEL030		70-kDa heat shock protein
1.9		YHR064		Heat shock protein
2.6	1.3	YBL008	HIR1	Histone transcription regulator
		YBL002	HTB2	Histone H2B.2
		YBL003	HTA2	Histone H2A.2
		YBR010	HHT1	Histone H3
3.3		YBR009	HHF1	Histone H4
3.9	2.4	YDR343	HXT6	High-affinity hexose transporter
	2.1	YHR092	HXT4	Moderate- to low-affinity glucose transporter
3.6		YAR071	PHO11	Secreted acid phosphatase, 56 kDa isozyme
	2.3	YLR096	KIN2	Ser/Thr protein kinase
2.5		YER102	RPS8B	Ribosomal protein S8.e
2.6		YBR181	RPS101	Ribosomal protein S6.e
2.6		YCR031	CRY1	40S ribosomal protein S14.e
2.7		YLR441	RP10A	Ribosomal protein S3.a.e
2.8		YHR141	RPL41B	Ribosomal protein L36a.e
2.8		YBL072	RPS8A	Ribosomal protein S8.e
2.8		YHL015	URP2	Ribosomal protein
2.8		YBR191	URP1	Ribosomal protein L21.e
3.1		YLR340	RPLA0	Acidic Ribosomal protein L10.e
3.3		YGL123	SUP44	Ribosomal protein
	5.8	YLR194		Hypothetical protein

500–1,000 arrays. Thus, a single primer pair provides enough starting material for up to  $\approx 50,000$  arrays.

Primers were synthesized to amplify yeast ORFs. Primer synthesis had a failure rate of  $<1\%$  in over 18 plates of synthesis as determined by standard trityl analysis (6). The success rate of the PCR amplifications using the primer pairs was 94% based on agarose gel analysis of each PCR. The purified PCR products were used to generate arrays. Two versions of the arrays were created for the experimental results presented here. The first array contained 2,287 elements and the second array batch contained 2,479 elements.

**Genome Arrays.** The amplified ORFs were arrayed onto glass at a spacing of 345 microns (Fig. 1). The high-density spacing of DNA samples allows the hybridization volumes to be minimized—volumes are a maximum of 10  $\mu$ l. The labeled probe can thus be maintained at relatively high concentrations, making 1–2  $\mu$ g of mRNA sufficient for analysis. This also obviates the need for a subsequent amplification step and thus avoids the risk of altering the relative ratios of different cDNA species in the sample.

**Genetic Analysis: Genomic Comparison of Unrelated Strains.** Microarrays allow efficient comparison of the genomes of different strains. Genomic DNA from Y55, an *S. cerevisiae* strain divergent from the reference strain S288c, was randomly labeled with Cy3-dUTP and hybridized simultaneously with the S288c DNA labeled with Cy5-dUTP. When a comparison between the hybridization of the DNA from the two strains was done, several

elements gave relatively little or no signal above background from the Cy3 channel (data not shown). These include SGE1, ASP3A-D, YLR156, YLR159, YLR161, ENA2 (YDR039 is ENA2), and YCR105. These results imply that the regions containing these genes are extremely divergent, or all together deleted from the strain. Subsequent attempts to generate PCR products from SGE1, ENA2, and ASP3A using Y55 DNA failed. This result supports the conclusion that these genes are likely to be missing from the Y55 genome. It is interesting to note that at least two of the regions absent in the Y55 genome have been previously shown or suggested to be deleted in mutant laboratory strains (14–16). In particular, the Asp-3 region appears to be highly prone to being deleted (15, 16).

These results indicate that gene arrays can be used to efficiently screen different strains of an organism for large deletion polymorphisms. A single hybridization and scan will reveal differences based on differential hybridization to particular elements. It is reasonable to suppose that an equivalent number of genes are present in the Y55 genome and absent in the S288c genome. This result should be viewed as a minimum estimate of the deletion polymorphisms that exist between these two unrelated strains as intergenic deletions or small intragenic deletions would not be detected because considerable hybridizing material would be remain. Sequence polymorphisms, such as deletions, are present in populations of every species and must at some level affect phenotype. One of the challenges of the genome era will be to critically examine sequence polymorphisms that exist in the natural gene pool relative to the reference genome sequence.






FIG. 2. ORF categories displaying differential expression between heat shocked and untreated cultures. Bars within categories correspond to individual ORFs. Green shaded bars correspond to relative increases in ORF expression under 25°C growth conditions. Red shaded bars correspond to relative increases in ORF expression under 39°C growth conditions.

**Gene Expression Analysis.** The arrays were used to examine gene expression in yeast grown under a variety of different conditions. Expression analysis is an ideal application of these arrays because a single hybridization provides quantitative expres-

sion data for thousands of genes. To better understand results for genes of known function, ORFs were placed in biologically relevant categories on the basis of function (e.g., amino acid catabolic genes) and/or pathways (e.g., the histidine biosynthesis pathway).

Table 2. Cold shock vs. control expression data

Ratio of gene expression		ORF	Gene	Description
Control	Cold			
	3.3	YOR153	PDR5	Pleiotropic drug resistance protein
2.4		YCR012	PGK1	Phosphoglycerate kinase
2.9		YCL040	GLK1	Aldohexose specific glucokinase
	1.4	YHR064		Heat shock protein
2.0		YJL034	KAR2	Nuclear fusion protein
2.1		YDR258	HSP78	Mitochondrial heat shock protein of clpb family of ATP-dependent proteases
2.2		YLL039	UBI4	Ubiquitin precursor
2.7		YLL026	HSP104	Heat shock protein
3.1		YER103	SSA4	Heat shock protein
3.3		YBR126	TPS1	$\alpha$ , $\alpha$ -Trehalose-phosphate synthase (UDP-forming)
3.8		YPL240	HSP82	Heat shock protein
7.9		YBR054	YRO2	Similarity to HSP30 heat shock protein Yro1p
7.9		YBR072	HSP26	Heat shock protein
16.5		YCR021	HSP30	Heat shock protein
1.8		YDR343	HXT6	High-affinity hexose transporter
2.1		YHR096	HXT5	Putative hexose transporter
2.4		YFR053	HXX1	Hexokinase I
2.8		YHR092	HXT4	Moderate- to low-affinity glucose transporter
3.4		YHR094	HXT1	Low-affinity hexose (glucose) transporter
	2.3	YHR089	GAR1	Nucleolar rRNA processing protein
	1.7	YLR048	NAB1B	40S ribosomal protein p40 homolog b
	1.7	YLR441	RP10A	Ribosomal protein S3a.e
	1.7	YLL045	RPL4B	Ribosomal protein L7a.e.B
	1.6	YLR029	RPL13A	Ribosomal protein L15.e
	1.6	YGL123	SUP44	Ribosomal protein
	3.1	YBR067	TIP1	Cold- and heat-shock-induced protein of the Srp1/Tip1p family
	2.2	YER011	TIR1	Cold-shock-induced protein of the Tir1p, Tip1p family
	2.0	YCR058		Hypothetical protein
	4.2	YKL102		Hypothetical protein

Table 3. Glucose vs. galactose expression data

Ratio of gene expression		ORF	Gene	Description
Glucose	Galactose			
2.1		YHR018	ARG4	Arginosuccinate lyase
3.5		YPR035	GLN1	Glutamate-ammonia ligase
2.8		YML116	ATR1	Aminotriazole and 4-nitroquinoline resistance protein
2.0		YMR303	ADH2	Alcohol dehydrogenase II
3.7		YBR145	ADH5	Alcohol dehydrogenase V
	3.2	YBL030	AAC2	ADP, ATP carrier protein 2
	2.9	YBR085	AAC3	ADP, ATP carrier protein
	2.7	YDR298	ATP5	H <sup>+</sup> -transporting ATP synthase $\delta$ chain precursor
	2.5	YBR039	ATP3	H <sup>+</sup> -transporting ATP synthase $\gamma$ chain precursor
	5.5	YML054	CYB2	Lactate dehydrogenase cytochrome <i>b</i> 2
	3.4	YML054	CYB2	Lactate dehydrogenase cytochrome <i>b</i> 2
	2.3	YKL150	MCR1	Cytochrome- <i>b</i> 5 reductase
	4.2	YBL045	COR1	Ubiquinol-cytochrome <i>c</i> reductase 44K core protein
	3.5	YDL067	COX9	Cytochrome <i>c</i> oxidase chain VIIA
	2.7	YLR038	COX12	Cytochrome <i>c</i> oxidase, subunit VIB
	2.6	YHR051	COX6	Cytochrome <i>c</i> oxidase subunit VI
	2.4	YLR395	COX8	Cytochrome <i>c</i> oxidase chain VIII
	2.3	YFR033	QCR6	Ubiquinol-cytochrome <i>c</i> reductase 17K protein
	23.7	YLR081	GAL2	Galactose (and glucose) permease
	21.9	YBR018	GAL7	UDP-glucose-hexose-1-phosphate uridylyltransferase
	21.8	YBR020	GAL1	Galactokinase
	19.5	YBR019	GAL10	UDP-glucose 4-epimerase
	14.7	YLR081	GAL2	Galactose (and glucose) permease
	8.6	YDR009	GAL3	Galactokinase
	3.0	YML051	GAL80(1)	Negative regulator for expression of galactose-induced genes
	2.8	YML051	GAL80(2)	Negative regulator for expression of galactose-induced genes
2.7		YER055	HIS1	ATP phosphoribosyltransferase
3.4		YBR248	HIS7	Glutamine amidotransferase/cyclase
				Phosphoribosyl-AMP cyclohydrolase/phosphoribosyl-ATP pyrophosphatase/histidinol dehydrogenase
7.4		YCL030	HIS4	
5.8		YKR080	MTD1	Methylenetetrahydrofolate dehydrogenase (NAD <sup>+</sup> )
6.0		YDR019	GCV1	Glycine decarboxylase T subunit
6.1		YLR058	SHM2	Serine hydroxymethyltransferase
	8.1	YML123	PHO84	High-affinity inorganic phosphate/H <sup>+</sup> symporter
3.5		YDR408	ADE8	Phosphoribosylglycinamide formyltransferase (GART)
3.6		YDR408	ADE8	Phosphoribosylglycinamide formyltransferase (GART)
4.4		YAR015	ADE1	Phosphoribosylamidoimidazole-succinocarboxamide synthase
5.6		YMR300	ADE4	Amidophosphoribosyltransferase
5.6		YOR128	ADE2	Phosphoribosylaminoimidazole carboxylase
6.0		YGL234	ADE5,7	Phosphoribosylamine-glycine ligase and phosphoribosylformylglycinamide cyclo-ligase
	6.3	YBL015	ACH1	Acetyl-CoA hydrolase

**Heat Shock Results.** A log phase culture growing in YEP/dextrose medium at 25°C was split in half. One half of the culture remained at 25°C whereas the other half of the culture was shifted to 39°C. mRNA was isolated from both cultures 1 h after heat shock for comparison on microarrays and, although this time point is not optimal for measuring induction of heat shock mRNAs (17), many known heat shock genes exhibited considerable induction at this time point (Table 1; Fig. 2). Down-regulation of genes in the ribosomal protein and histone gene categories was also observed. Differential expression between the heat-shocked culture and the control was also observed for many other genes. Genes in many categories, such as amino acid catabolism and amino acid synthesis, exhibited a mixed response with some genes showing little or no differential expression and other genes showing a significant increase or decrease in gene expression in response to heat shock (Table 1; Fig. 2).

**Cold Shock Results.** A log phase culture growing in YEP/dextrose medium at 37°C was split in half. One half of the culture remained at 37°C while the other half of the culture was shifted to 18°C. mRNA was isolated from both cultures 1 h after cold shock for comparison on microarrays. As expected,

two known cold shock genes (TIP1, TIR1) were expressed at a significantly higher level in the cold-shocked culture. Genes in other functional categories, such as glucose metabolism and heat shock displayed a mixed response with expression of some genes being unaffected and other genes exhibiting significant up- or down-regulation in response to cold shock (Table 2).

**Steady-State Galactose vs. Glucose Results.** mRNA was isolated from steady-state log phase YEP galactose and YEP glucose grown cultures for comparison on the microarrays. As expected, the GAL genes were expressed at a much higher level in the galactose culture. Many genes were differentially expressed in these cultures that were not *a priori* expected to exhibit differential expression. For example, some genes in the amino acid catabolic category were up-regulated in the galactose culture whereas genes in the one-carbon metabolism and purine categories were largely or entirely down-regulated in the galactose culture (Table 3). Genes in other categories, such as amino acid synthesis, abc transporter, cytochrome *c*, and cytochrome *b*, exhibited mixed responses; some genes in a category showed little or no obvious differential expression whereas other genes in the same category showed significant differential expression in the galactose and glucose cultures.

## DISCUSSION

The results of these experiments show that many genes are differentially expressed under the three environmental conditions described here. The expected and predicted changes in gene expression, such as HSP12 in the heat-shocked culture, TIP1 in the cold-shocked culture, and GAL2 in the steady-state galactose culture, were observed in every case. However, in addition to the expected changes in gene expression, significant differential expression was also observed for many other genes that would not, *a priori*, be expected to be differentially expressed. For example, expression of PHO11 decreased and expression of YLR194, KIN2, and HXT6 increased in the heat shocked culture. Expression of MST1 and APE3 decreased and expression of PDR5 and GAR1 increased in the cold-shocked culture. In addition, ADE4 and SER2 were expressed at reduced levels whereas PHO84 and ACH1 were expressed at higher levels in cells grown in galactose compared with cells grown in glucose. Differential expression of these and many other genes was specific to one of these three environmental conditions.

Many other genes were found to be differentially expressed under more than one condition. When differentially expressed genes in cold- and heat-shocked cultures were compared, 30 genes were found in common. Of these 30 genes, 28 showed inverse expression (i.e., increased expression under one condition and decreased expression under the other condition). Two genes, YCR058 and YKL102, showed elevated expression in response to both cold and heat shock. Fifteen genes were found to be differentially expressed in both the heat-shocked and steady-state galactose cultures: 9 genes showed increased expression and 5 showed decreased expression under both conditions. Twenty genes were differentially expressed in both the cold-shocked and steady-state galactose cultures: 8 genes showed decreased expression and 5 genes showed increased expression under both conditions. Six genes showed increased expression in the galactose culture and decreased expression in the cold shocked culture. One gene (ODP1) showed increased expression in both the cold-shocked and steady-state galactose cultures.

Gene expression is affected in a global fashion when environmental conditions are changed and both expected and unexpected genes are affected. There is also overlap in the genes that are differentially expressed under quite different environmental conditions. These results can be rationalized by considering the high degree of cross-pathway regulation in yeast. For example, there is evidence for cross-pathway regulation between (i) carbon and nitrogen metabolism (18), (ii) phosphate and sulfate metabolism (19), and (iii) purine, phosphate, and amino acid metabolism (20–24). There are also examples of the interaction of general and specific transcription factors (25, 26). Finally, within the broad class of amino acid biosynthetic genes, there is evidence for amino acid specific regulation of some genes, regulation via general control for other genes, and regulation via both specific and general control for other genes (22, 27–30).

Cross-pathway regulation arises from the complex structure of promoters. Virtually all promoters contain sites for multiple transcription factors and, therefore, virtually all genes are subject to combinatorial regulation. For example, the HIS4 promoter contains binding sites for GCN4 (the general amino acid control transcription factor), PHO2/BAS2 (a transcriptional regulator of phosphatase and purine biosynthetic genes), and BAS1 (a transcriptional regulator of purine biosynthetic genes) (31). It is likely that the complex effects on gene expression described in this work are a direct consequence of the combinatorial regulation of gene expression.

These findings illustrate the power of the highly parallel whole genome approach when examining gene expression. The global effects of environmental change on gene expression can now be directly visualized. It is clear that determining the mechanism(s) and the functional role of the dramatic global effects on gene

expression in different environments will be a significant challenge. The era of whole genome analysis will, ultimately, allow researchers to switch from the very focused single gene/promoter view of gene expression and instead view the cell more as a large complex network of gene regulatory pathways.

With the entire sequence of this model organism known, new approaches have been developed that allow for genome wide analyses (32, 33) of gene function. The genome microarrays represent a novel tool for genetic and expression analysis of the yeast genome. This pilot study uses arrays containing >35% of the yeast ORFs and it is clear that the entire set of ORFs from the yeast genome can be arrayed using the directed primer based strategy detailed here. Recent advances in arraying technology will allow all 6,100 ORFs to be arrayed in an area of less than 1.8 cm<sup>2</sup>. Furthermore, as the technology improves, detection limits will allow less than 500 ng of starting mRNA material to be used for making probe.

The genome arrays provide for a robust, fully automated approach toward examining genome structure and gene function. They allow for comparisons between different genomes as well as a detailed study of gene expression at the global level. This research will help to elucidate relationships between genes and allow the researcher to understand gene function by understanding expression patterns across the yeast genome.

Support was provided by National Institutes of Health Grant P01HG00205.

1. Fleischmann, R. D., Adams, M. D., White, O., Clayton, R. A., Kirkness, E. F., *et al.* (1995) *Science* 269, 496–512.
2. Fraser, C. M., Gocayne, J. D., White, O., Adams, M. D., Clayton, R. A., *et al.* (1995) *Science* 270, 397–403.
3. Bult, C. J., White, O., Olsen, G. J., Zhou, L., Fleischmann, R. D., *et al.* (1996) *Science* 273, 1058–1073.
4. Sulston, J., Du, Z., Thomas, K., Wilson, R., Hillier, L., *et al.* (1992) *Nature (London)* 356, 37.
5. Newman, T., de Bruijn, F. J., Green, P., Keegstra, K., Kende, H., *et al.* (1994) *Plant Physiol.* 106, 1241–1255.
6. Lashkari, D. A., Hunicke-Smith, S. P., Norgren, R. M., Davis, R. W. & Brennan, T. (1995) *Proc. Natl. Acad. Sci. USA* 92, 7912–7915.
7. Schena, M., Shalon, D., Davis, R. W. & Brown, P. O. (1995) *Science* 270, 467–470.
8. Shalon, D., Smith, S. & Brown, P. O. (1996) *Genome Res.* 6, 639–645.
9. Heller, R. A., Schena, M., Chai, A., Shalon, D., Bedilion, T., Gilmore, J., Woolley, D. E. & Davis, R. W. (1997) *Proc. Natl. Acad. Sci. USA* 94, 2150–2155.
10. DeRisi, J., Penland, L., Brown, P. O., Bittner, M. L., Meltzer, P. S., Ray, M., Chen, Y., Su, Y. & Trent, J. M. (1996) *Nat. Genet.* 14, 457–460.
11. Nelson, S. F., McCusker, J. H., Sander, M. A., Kee, Y., Modrich, P. & Brown P. O. (1993) *Nat. Genet.* 4, 11–18.
12. Hoffman, C. S. & Winston, F. (1989) *Gene* 84, 473–479.
13. Schmitt, M., Brown, T. & Trumpower, B. (1990) *Nucleic Acids Res.* 18, 3091.
14. Ehrenhofer-Murray, A. E., Wurgler, F. E. & Sengstag, C. (1994) *Mol. Gen. Genet.* 244, 287–294.
15. Kim, K.-W., Kamerud, J. Q., Livingston, D. M. & Roon, R. J. (1988) *J. Biol. Chem.* 263, 11948–11953.
16. Kim, K.-W. & Roon, R. J. (1984) *J. Bacteriol.* 157, 958–961.
17. Craig, E. A. (1992) in *The Molecular Biology of the Yeast Saccharomyces: Gene Expression*, eds. Jones, E. W., Pringle, J. R. & Broach, J. R. (Cold Spring Harbor Lab. Press, Plainview, NY), Vol. 2, pp. 501–537.
18. Dang, V. D., Bohn, C., Bolotin-Fukuhara, M. & Daignan-Fornier, B. (1996) *J. Bacteriol.* 178, 1842–1849.
19. O'Connell, K. F. & Baker, R. E. (1992) *Genetics* 132, 63–73.
20. Braus, G., Mosch, H. U., Vogel, K., Hinnen, A. & Hutter, R. (1989) *EMBO J.* 8, 939–945.
21. Mosch, H. U., Scheier, B., Lahti, R., Mantsala, P. & Braus, G. H. (1991) *J. Biol. Chem.* 266, 20453–20456.
22. Mitchell, A. P. & Magasanik, B. (1984) *Mol. Cell. Biol.* 4, 2767–2773.
23. Daignan-Fornier, B. & Fink, G. R. (1992) *Proc. Natl. Acad. Sci. USA* 89, 6746–6750.
24. Tice-Baldwin, K., Fink, G. R. & Arndt, K. T. (1989) *Science* 246, 931–935.
25. Messenguy, F. & Dubois, E. (1993) *Mol. Cell. Biol.* 13, 2586–2592.
26. Devlin, C., Tice-Baldwin, K., Shore, D. & Arndt, K. T. (1991) *Mol. Cell. Biol.* 11, 3642–3651.
27. Magasanik, B. (1992) in *The Molecular and Cellular Biology of the Yeast Saccharomyces: Gene Expression*, eds. Jones, E. W., Pringle, J. R. & Broach, J. R. (Cold Spring Harbor Lab. Press, Plainview, NY), Vol. 2, pp. 283–317.
28. Hinnebusch, A. G. (1992) in *The Molecular and Cellular Biology of the Yeast Saccharomyces: Gene Expression*, eds. Jones, E. W., Pringle, J. R. & Broach, J. R. (Cold Spring Harbor Lab. Press, Plainview, NY), Vol. 2, pp. 319–414.
29. Brisco, P. R. & Kohlhaw, G. B. (1990) *J. Biol. Chem.* 265, 11667–11675.
30. O'Connell, K. F., Surdin-Kerjan, Y. & Baker, R. E. (1995) *Mol. Cell. Biol.* 15, 1879–1888.
31. Arndt, K. T., Styles, C. & Fink, G. R. (1987) *Science* 237, 874–880.
32. Smith, V., Chou, K. N., Lashkari, D., Botstein, D. & Brown, P. O. (1996) *Science* 274, 2069–2074.
33. Shoemaker, D. D., Lashkari, D. A., Morris, D., Mittman, M. & Davis, R. W. (1996) *Nat. Genet.* 14, 450–456.



- Fischer-Vize, *Science* 270, 1828 (1995).
35. T. C. James and S. C. Elgin, *Mol. Cell Biol.* 6, 3862 (1986); R. Paro and D. S. Hogness, *Proc. Natl. Acad. Sci. U.S.A.* 88, 263 (1991); B. Tschiersch et al., *EMBO J.* 13, 3822 (1994); M. T. Madireddi et al., *Cell* 87, 75 (1996); D. G. Stokes, K. D. Tartof, R. P. Perry, *Proc. Natl. Acad. Sci. U.S.A.* 93, 7137 (1996).
36. P. M. Palosaari et al., *J. Biol. Chem.* 266, 10750 (1991); A. Schmitz, K. H. Gartemann, J. Fiedler, E. Grund, R. Eichenlaub, *Appl. Environ. Microbiol.* 58, 4068 (1992); V. Sharma, K. Suvama, R. Meganathan, M. E. Hudspeth, *J. Bacteriol.* 174, 5057 (1992); M. Kanazawa et al., *Enzyme Protein* 47, 9 (1993); Z. L. Boynton, G. N. Bennet, F. B. Rudolph, *J. Bacteriol.* 178, 3015 (1996).
37. M. Ho et al., *Cell* 77, 869 (1994).
38. W. Hendriks et al., *J. Cell Biochem.* 59, 418 (1995).
39. We thank H. Skaletsky and F. Lewitter for help with

sequence analysis; Lawrence Livermore National Laboratory for the flow-sorted Y cosmid library; and P. Bain, A. Bortvin, A. de la Chapelle, G. Fink, K. Jegalian, T. Kawaguchi, E. Lander, H. Lodish, P. Matsudaira, D. Menke, U. RajBhandary, R. Reijo, S. Rozen, A. Schwartz, C. Sun, and C. Tifford for comments on the manuscript. Supported by NIH.

28 April 1997; accepted 9 September 1997

## Exploring the Metabolic and Genetic Control of Gene Expression on a Genomic Scale

Joseph L. DeRisi, Vishwanath R. Iyer, Patrick O. Brown\*

DNA microarrays containing virtually every gene of *Saccharomyces cerevisiae* were used to carry out a comprehensive investigation of the temporal program of gene expression accompanying the metabolic shift from fermentation to respiration. The expression profiles observed for genes with known metabolic functions pointed to features of the metabolic reprogramming that occur during the diauxic shift, and the expression patterns of many previously uncharacterized genes provided clues to their possible functions. The same DNA microarrays were also used to identify genes whose expression was affected by deletion of the transcriptional co-repressor *TUP1* or overexpression of the transcriptional activator *YAP1*. These results demonstrate the feasibility and utility of this approach to genomewide exploration of gene expression patterns.

The complete sequences of nearly a dozen microbial genomes are known, and in the next several years we expect to know the complete genome sequences of several metazoans, including the human genome. Defining the role of each gene in these genomes will be a formidable task, and understanding how the genome functions as a whole in the complex natural history of a living organism presents an even greater challenge.

Knowing when and where a gene is expressed often provides a strong clue as to its biological role. Conversely, the pattern of genes expressed in a cell can provide detailed information about its state. Although regulation of protein abundance in a cell is by no means accomplished solely by regulation of mRNA, virtually all differences in cell type or state are correlated with changes in the mRNA levels of many genes. This is fortuitous because the only specific reagent required to measure the abundance of the mRNA for a specific gene is a cDNA sequence. DNA microarrays, consisting of thousands of individual gene sequences printed in a high-density array on a glass microscope slide (1, 2), provide a practical and economical tool for studying gene expression on a very large scale (3–6).

*Saccharomyces cerevisiae* is an especially

favorable organism in which to conduct a systematic investigation of gene expression. The genes are easy to recognize in the genome sequence, cis regulatory elements are generally compact and close to the transcription units, much is already known about its genetic regulatory mechanisms, and a powerful set of tools is available for its analysis.

A recurring cycle in the natural history of yeast involves a shift from anaerobic (fermentation) to aerobic (respiration) metabolism. Inoculation of yeast into a medium rich in sugar is followed by rapid growth fueled by fermentation, with the production of ethanol. When the fermentable sugar is exhausted, the yeast cells turn to ethanol as a carbon source for aerobic growth. This switch from anaerobic growth to aerobic respiration upon depletion of glucose, referred to as the diauxic shift, is correlated with widespread changes in the expression of genes involved in fundamental cellular processes such as carbon metabolism, protein synthesis, and carbohydrate storage (7). We used DNA microarrays to characterize the changes in gene expression that take place during this process for nearly the entire genome, and to investigate the genetic circuitry that regulates and executes this program.

Yeast open reading frames (ORFs) were amplified by the polymerase chain reaction (PCR), with a commercially available set of primer pairs (8). DNA microarrays, containing approximately 6400 distinct DNA sequences, were printed onto glass slides by

using a simple robotic printing device (9). Cells from an exponentially growing culture of yeast were inoculated into fresh medium and grown at 30°C for 21 hours. After an initial 9 hours of growth, samples were harvested at seven successive 2-hour intervals, and mRNA was isolated (10). Fluorescently labeled cDNA was prepared by reverse transcription in the presence of Cy3(green)- or Cy5(red)-labeled deoxyuridine triphosphate (dUTP) (11) and then hybridized to the microarrays (12). To maximize the reliability with which changes in expression levels could be discerned, we labeled cDNA prepared from cells at each successive time point with Cy5, then mixed it with a Cy3-labeled "reference" cDNA sample prepared from cells harvested at the first interval after inoculation. In this experimental design, the relative fluorescence intensity measured for the Cy3 and Cy5 fluoros at each array element provides a reliable measure of the relative abundance of the corresponding mRNA in the two cell populations (Fig. 1). Data from the series of seven samples (Fig. 2), consisting of more than 43,000 expression-ratio measurements, were organized into a database to facilitate efficient exploration and analysis of the results. This database is publicly available on the Internet (13).

During exponential growth in glucose-rich medium, the global pattern of gene expression was remarkably stable. Indeed, when gene expression patterns between the first two cell samples (harvested at a 2-hour interval) were compared, mRNA levels differed by a factor of 2 or more for only 19 genes (0.3%), and the largest of these differences was only 2.7-fold (14). However, as glucose was progressively depleted from the growth media during the course of the experiment, a marked change was seen in the global pattern of gene expression. mRNA levels for approximately 710 genes were induced by a factor of at least 2, and the mRNA levels for approximately 1030 genes declined by a factor of at least 2. Messenger RNA levels for 183 genes increased by a factor of at least 4, and mRNA levels for 203 genes diminished by a factor of at least 4. About half of these differentially expressed genes have no currently recognized function and are not yet named. Indeed, more than 400 of the differentially expressed genes have no apparent homology

Department of Biochemistry, Stanford University School of Medicine, Howard Hughes Medical Institute, Stanford, CA 94305–5428, USA.

\*To whom correspondence should be addressed. E-mail: pbrown@crgm.stanford.edu



to any gene whose function is known (15). The responses of these previously uncharacterized genes to the diauxic shift therefore provides the first small clue to their possible roles.

The global view of changes in expression of genes with known functions provides a vivid picture of the way in which the cell adapts to a changing environment. Figure 3 shows a portion of the yeast metabolic pathways involved in carbon and energy metabolism. Mapping the changes we observed in the mRNAs encoding each enzyme onto this framework allowed us to infer the redirection in the flow of metabolites through this system. We observed large inductions of the genes coding for the enzymes aldehyde dehydrogenase (ALD2) and acetyl-coenzyme A (CoA) synthase (ACS1), which function together to convert the products of alcohol dehydrogenase into acetyl-CoA, which in turn is used to fuel the tricarboxylic acid (TCA) cycle and the glyoxylate cycle. The concomitant shutdown of transcription of the genes encoding pyruvate decarboxylase and induction of pyruvate carboxylase rechannels pyruvate away from acetaldehyde, and instead to oxalacetate, where it can serve to supply the TCA cycle and gluconeogenesis. Induction of the pivotal genes *PCK1*, encoding phosphoenolpyruvate carboxykinase, and *FBP1*, encoding fructose 1,6-bisphosphatase, switches the directions of two key irreversible steps in glycolysis, reversing the flow of metabolites along the reversible steps of the glycolytic pathway toward the essential biosynthetic precursor, glucose-6-phosphate. Induction of the genes coding for the trehalose synthase and glycogen synthase complexes promotes channeling of glucose-6-phosphate into these carbohydrate storage pathways.


Just as the changes in expression of genes encoding pivotal enzymes can provide insight into metabolic reprogramming, the behavior of large groups of functionally related genes can provide a broad view of the systematic way in which the yeast cell adapts to a changing environment (Fig. 4). Several classes of genes, such as cytochrome c-related genes and those involved in the TCA/glyoxylate cycle and carbohydrate storage, were coordinately induced by glucose exhaustion. In contrast, genes devoted to protein synthesis, including ribosomal proteins, tRNA synthetases, and translation, elongation, and initiation factors, exhibited a coordinated decrease in expression. More than 95% of ribosomal genes showed at least twofold decreases in expression during the diauxic shift (Fig. 4) (13). A noteworthy and illuminating exception was that the

genes encoding mitochondrial ribosomal genes were generally induced rather than repressed after glucose limitation, highlighting the requirement for mitochondrial biogenesis (13). As more is learned about the functions of every gene in the yeast genome, the ability to gain insight into a cell's response to a changing environment through its global gene expression patterns will become increasingly powerful.

Several distinct temporal patterns of expression could be recognized, and sets of genes could be grouped on the basis of the similarities in their expression patterns. The characterized members of each of these groups also shared important similarities in their functions. Moreover, in most cases, common regulatory mechanisms could be inferred for sets of genes with similar expression profiles. For example, seven genes showed a late induction profile, with mRNA levels increasing by more than ninefold at

the last timepoint but less than threefold at the preceding timepoint (Fig. 5B). All of these genes were known to be glucose-repressed, and five of the seven were previously noted to share a common upstream activating sequence (UAS), the carbon source response element (CSRE) (16–20). A search in the promoter regions of the remaining two genes, *ACR1* and *IDP2*, revealed that *ACR1*, a gene essential for ACS1 activity, also possessed a consensus CSRE motif, but interestingly, *IDP2* did not. A search of the entire yeast genome sequence for the consensus CSRE motif revealed only four additional candidate genes, none of which showed a similar induction.

Examples from additional groups of genes that shared expression profiles are illustrated in Fig. 5, C through F. The sequences upstream of the named genes in Fig. 5C all contain stress response elements (STRE), and with the exception



**Fig. 1.** Yeast genome microarray. The actual size of the microarray is 18 mm by 18 mm. The microarray was printed as described (9). This image was obtained with the same fluorescent scanning confocal microscope used to collect all the data we report (49). A fluorescently labeled cDNA probe was prepared from mRNA isolated from cells harvested shortly after inoculation (culture density of  $<5 \times 10^6$  cells/ml and media glucose level of 19 g/liter) by reverse transcription in the presence of Cy3-dUTP. Similarly, a second probe was prepared from mRNA isolated from cells taken from the same culture 9.5 hours later (culture density of  $\sim 2 \times 10^8$  cells/ml, with a glucose level of  $<0.2$  g/liter) by reverse transcription in the presence of Cy5-dUTP. In this image, hybridization of the Cy3-dUTP-labeled cDNA (that is, mRNA expression at the initial timepoint) is represented as a green signal, and hybridization of Cy5-dUTP-labeled cDNA (that is, mRNA expression at 9.5 hours) is represented as a red signal. Thus, genes induced or repressed after the diauxic shift appear in this image as red and green spots, respectively. Genes expressed at roughly equal levels before and after the diauxic shift appear in this image as yellow spots.

of HSP42, have previously been shown to be controlled at least in part by these elements (21–24). Inspection of the sequences upstream of HSP42 and the two uncharacterized genes shown in Fig. 5C, YKL026c, a hypothetical protein with similarity to glutathione peroxidase, and YGR043c, a putative transaldolase, revealed that each of these genes also possess repeated upstream copies of the stress-responsive CCCCT motif. Of the 13 additional genes in the yeast genome that shared this expression profile [including HSP30, ALD2, OM45, and 10 uncharacterized ORFs (25)], nine contained one or more recognizable STRE sites in their upstream regions.

The heterotrimeric transcriptional activator complex HAP2,3,4 has been shown to be responsible for induction of several genes important for respiration (26–28). This complex binds a degenerate consensus sequence known as the CCAAT box (26). Computer analysis, using the consensus sequence TNRYTGGB (29), has suggested that a large number of genes involved in respiration may be specific targets of HAP2,3,4 (30). Indeed, a putative HAP2,3,4 binding site could be found in the sequences upstream of each of the seven cytochrome c-related genes that showed the greatest magnitude of induction (Fig. 5D). Of 12 additional cytochrome c-related genes that were induced, HAP2,3,4 binding sites were present in all but one. Significantly, we found that transcription of HAP4 itself was induced nearly ninefold concomitant with the diauxic shift.

Control of ribosomal protein biogenesis is mainly exerted at the transcriptional level, through the presence of a common upstream-activating element (UAS<sub>mp</sub>) that is recognized by the Rap1 DNA-binding protein (31, 32). The expression profiles of seven ribosomal proteins are shown in Fig. 5F. A search of the sequences upstream of all seven genes revealed consensus Rap1-binding motifs (33). It has been suggested that declining Rap1 levels in the cell during starvation may be responsible for the decline in ribosomal protein gene expression (34). Indeed, we observed that the abundance of RAP1 mRNA diminished by 4.4-fold, at about the time of glucose exhaustion.

Of the 149 genes that encode known or putative transcription factors, only two, HAP4 and SIP4, were induced by a factor of more than threefold at the diauxic shift. SIP4 encodes a DNA-binding transcriptional activator that has been shown to interact with Snf1, the “master regulator” of glucose repression (35). The eightfold induction of SIP4 upon depletion of glucose strongly suggests a role in the induction of


downstream genes at the diauxic shift.

Although most of the transcriptional responses that we observed were not previously known, the responses of many genes during the diauxic shift have been described. Comparison of the results we obtained by DNA microarray hybridization with previously reported results therefore provided a strong test of the sensitivity and accuracy of this approach. The expression patterns we observed for previously characterized genes showed almost perfect concordance with previously published results (36). Moreover, the differential expression measurements obtained by DNA microarray hybridization were reproducible in duplicate experiments. For example, the remarkable changes in gene expression between cells harvested immediately after inoculation and immediately after the diauxic shift (the first and sixth intervals in this time series) were measured in duplicate, independent DNA microarray hybridizations. The correlation coefficient for two complete sets of expression ratio measurements was 0.87, and for more than 95% of the genes, the expres-

sion ratios measured in these duplicate experiments differed by less than a factor of 2. However, in a few cases, there were discrepancies between our results and previous results, pointing to technical limitations that will need to be addressed as DNA microarray technology advances (37, 38). Despite the noted exceptions, the high concordance between the results we obtained in these experiments and those of previous studies provides confidence in the reliability and thoroughness of the survey.

The changes in gene expression during this diauxic shift are complex and involve integration of many kinds of information about the nutritional and metabolic state of the cell. The large number of genes whose expression is altered and the diversity of temporal expression profiles observed in this experiment highlight the challenge of understanding the underlying regulatory mechanisms. One approach to defining the contributions of individual regulatory genes to a complex program of this kind is to use DNA microarrays to identify genes whose expression is affected

**Fig. 2.** The section of the array indicated by the gray box in Fig. 1 is shown for each of the experiments described here. Representative genes are labeled. In each of the arrays used to analyze gene expression during the diauxic shift, red spots represent genes that were induced relative to the initial timepoint, and green spots represent genes that were repressed relative to the initial timepoint. In the arrays used to analyze the effects of the *tup1*Δ mutation and YAP1 overexpression, red spots represent genes whose expression was increased, and green spots represent genes whose expression was decreased by the genetic modification. Note that distinct sets of genes are induced and repressed in the different experiments. The complete images of each of these arrays can be viewed on the Internet (13). Cell density as measured by optical density (OD) at 600 nm was used to measure the growth of the culture.




by mutations in each putative regulatory gene. As a test of this strategy, we analyzed the genome-wide changes in gene expression that result from deletion of the *TUP1* gene. Transcriptional repression of many genes by glucose requires the DNA-binding repressor

Mig1 and is mediated by recruiting the transcriptional co-repressors Tup1 and Cyc8/Ssn6 (39). Tup1 has also been implicated in repression of oxygen-regulated, mating-type-specific, and DNA-damage-inducible genes (40).

Wild-type yeast cells and cells bearing a deletion of the *TUP1* gene (*tup1Δ*) were grown in parallel cultures in rich medium containing glucose as the carbon source. Messenger RNA was isolated from exponentially growing cells from the two populations and used to prepare cDNA labeled with Cy3 (green) and Cy5 (red), respectively (11). The labeled probes were mixed and simultaneously hybridized to the microarray. Red spots on the microarray therefore represented genes whose transcription was induced in the *tup1Δ* strain, and thus presumably repressed by Tup1 (41). A representative section of the microarray (Fig. 2, bottom middle panel) illustrates that the genes whose expression was affected by the *tup1Δ* mutation, were, in general, distinct from those induced upon glucose exhaustion [complete images of all the arrays shown in Fig. 2 are available on the Internet (13)]. Nevertheless, 34 (10%) of the genes that were induced by a factor of at least 2 after the diauxic shift were similarly induced by deletion of *TUP1*, suggesting that these genes may be subject to *TUP1*-mediated repression by glucose. For example, *SUC2*, the gene encoding invertase, and all five hexose transporter genes that were induced during the course of the diauxic shift were similarly induced, in duplicate experiments, by the deletion of *TUP1*.

The set of genes affected by Tup1 in this experiment also included  $\alpha$ -glucosidases, the mating-type-specific genes *MFA1* and *MFA2*, and the DNA damage-inducible *RNR2* and *RNR4*, as well as genes involved in flocculation and many genes of unknown function. The hybridization signal corresponding to expression of *TUP1* itself was also severely reduced because of the (incomplete) deletion of the transcription unit in the *tup1Δ* strain, providing a positive control in the experiment (42).

Many of the transcriptional targets of Tup1 fell into sets of genes with related biochemical functions. For instance, although only about 3% of all yeast genes appeared to be *TUP1*-repressed by a factor of more than 2 in duplicate experiments under these conditions, 6 of the 13 genes that have been implicated in flocculation (15) showed a reproducible increase in expression of at least twofold when *TUP1* was deleted. Another group of related genes that appeared to be subject to *TUP1* repression encodes the serine-rich cell wall mannoproteins, such as *Tip1* and *Tir1/Srp1* which are induced by cold shock and other stresses (43), and similar, serine-poor proteins, the seripauperins (44). Messenger RNA levels for 23 of the 26 genes in this group were reproducibly elevated by at least 2.5-fold in the *tup1Δ*



**Fig. 3.** Metabolic reprogramming inferred from global analysis of changes in gene expression. Only key metabolic intermediates are identified. The yeast genes encoding the enzymes that catalyze each step in this metabolic circuit are identified by name in the boxes. The genes encoding succinyl-CoA synthase and glycogen-debranching enzyme have not been explicitly identified, but the ORFs YGR244 and YPR184 show significant homology to known succinyl-CoA synthase and glycogen-debranching enzymes, respectively, and are therefore included in the corresponding steps in this figure. Green boxes with white lettering identify genes whose expression increases in the diauxic shift. The magnitude of induction or repression is indicated for these genes. For multimeric enzyme complexes, such as succinate dehydrogenase, the indicated fold-induction represents an unweighted average of all the genes listed in the box. Black and white boxes indicate no significant differential expression (less than twofold). The direction of the arrows connecting reversible enzymatic steps indicate the direction of the flow of metabolic intermediates, inferred from the gene expression pattern, after the diauxic shift. Arrows representing steps catalyzed by genes whose expression was strongly induced are highlighted in red. The broad gray arrows represent major increases in the flow of metabolites after the diauxic shift, inferred from the indicated changes in gene expression.



strain, and 18 of these genes were induced by more than sevenfold when *TUP1* was deleted. In contrast, none of 83 genes that could be classified as putative regulators of the cell division cycle were induced more than twofold by deletion of *TUP1*. Thus, despite the diversity of the regulatory systems that employ Tup1, most of the genes that it regulates under these conditions fall into a limited number of distinct functional classes.

Because the microarray allows us to monitor expression of nearly every gene in yeast, we can, in principle, use this approach to identify all the transcriptional targets of a regulatory protein like Tup1. It is important to note, however, that in any single experiment of this kind we can only recognize those target genes that are normally repressed (or induced) under the conditions of the experiment. For instance, the experiment described here analyzed a MAT  $\alpha$  strain in which *MFA1* and *MFA2*, the genes encoding the  $\alpha$ -factor mating pheromone precursor, are normally repressed. In the isogenic *tup1* $\Delta$  strain, these genes were inappropriately expressed, reflecting the role that Tup1 plays in their repression. Had we instead carried out this experiment with a MAT  $\alpha$  strain (in which expression of *MFA1* and *MFA2* is not repressed), it would not have been possible to conclude anything regarding the role of Tup1 in the repression of these genes. Conversely, we cannot distinguish indirect effects of the chronic absence of Tup1 in the mutant strain from effects directly attributable to its participation in repressing the transcription of a gene.

Another simple route to modulating the activity of a regulatory factor is to overexpress the gene that encodes it. *YAP1* encodes a DNA-binding transcription factor belonging to the b-zip class of DNA-binding proteins. Overexpression of *YAP1* in yeast confers increased resistance to hydrogen peroxide, *o*-phenanthroline, heavy metals, and osmotic stress (45). We analyzed differential gene expression between a wild-type strain bearing a control plasmid and a strain with a plasmid expressing *YAP1* under the control of the strong *GALI-10* promoter, both grown in galactose (that is, a condition that induces *YAP1* overexpression). Complementary DNA from the control and *YAP1* overexpressing strains, labeled with Cy3 and Cy5, respectively, was prepared from mRNA isolated from the two strains and hybridized to the microarray. Thus, red spots on the array represent genes that were induced in the strain overexpressing *YAP1*.


Of the 17 genes whose mRNA levels increased by more than threefold when

*YAP1* was overexpressed in this way, five bear homology to aryl-alcohol oxidoreductases (Fig. 2 and Table 1). An additional four of the genes in this set also belong to the general class of dehydrogenases/oxidoreductases. Very little is known about the role of aryl-alcohol oxidoreductases in *S. cerevisiae*, but these enzymes have been isolated from ligninolytic fungi, in which they participate in coupled redox reactions, oxidizing aromatic, and aliphatic unsaturated alcohols to aldehydes with the production of hydrogen peroxide (46, 47). The fact that a remarkable fraction of the targets identified in this experiment belong to the same small, functional group of oxidoreductases suggests that these genes

might play an important protective role during oxidative stress. Transcription of a small number of genes was reduced in the strain overexpressing Yap1. Interestingly, many of these genes encode sugar permeases or enzymes involved in inositol metabolism.

We searched for Yap1-binding sites (TTACTAA or TGAATA) in the sequences upstream of the target genes we identified (48). About two-thirds of the genes that were induced by more than threefold upon Yap1 overexpression had one or more binding sites within 600 bases upstream of the start codon (Table 1), suggesting that they are directly regulated by Yap1. The absence of canonical Yap1-bind-

**Fig. 4.** Coordinated regulation of functionally related genes. The curves represent the average induction or repression ratios for all the genes in each indicated group. The total number of genes in each group was as follows: ribosomal proteins, 112; translation elongation and initiation factors, 25; tRNA synthetases (excluding mitochondrial synthetases), 17; glycogen and trehalose synthesis and degradation, 15; cytochrome c oxidase and reductase proteins, 19; and TCA- and glyoxylate-cycle enzymes, 24.



**Table 1.** Genes induced by *YAP1* overexpression. This list includes all the genes for which mRNA levels increased by more than twofold upon *YAP1* overexpression in both of two duplicate experiments, and for which the average increase in mRNA level in the two experiments was greater than threefold (50). Positions of the canonical Yap1 binding sites upstream of the start codon, when present, and the average fold-increase in mRNA levels measured in the two experiments are indicated.

ORF	Distance of Yap1 site from ATG	Gene	Description	Fold-increase
YNL331C	162-222 (5 sites)	<i>YAP1</i>	Putative aryl-alcohol reductase	12.9
YKL071W			Similarity to bacterial <i>csgA</i> protein	10.4
YML007W			Transcriptional activator involved in oxidative stress response	9.8
YFL056C	223, 242		Homology to aryl-alcohol dehydrogenases	9.0
YLL060C	98		Putative glutathione transferase	7.4
YOL165C	266		Putative aryl-alcohol dehydrogenase (NADP+)	7.0
YCR107W	409	<i>ATR1</i>	Putative aryl-alcohol reductase	6.5
YML116W			Aminotriazole and 4-nitroquinoline resistance protein	6.5
YBR008C			Homology to benomyl/methotrexate resistance protein	6.1
YCLX08C	148, 212	<i>OYE3</i>	Hypothetical protein	6.1
YJR155W			Putative aryl-alcohol dehydrogenase	6.0
YPL171C			NAPDH dehydrogenase (old yellow enzyme), isoform 3	5.8
YLR460C	167, 317		Homology to hypothetical proteins YCR102c and YNL134c	4.7
YKR076W	178		Homology to hypothetical protein YMR251w	4.5
YHR179W	327	<i>OYE2</i>	NAD(P)H oxidoreductase (old yellow enzyme), isoform 1	4.1
YML131W	507		Similarity to <i>A. thaliana</i> zeta-crystallin homolog	3.7
YOL126C		<i>MDH2</i>	Malate dehydrogenase	3.3

ing sites upstream of the others may reflect an ability of Yap1 to bind sites that differ from the canonical binding sites, perhaps in cooperation with other factors, or less likely, may represent an indirect effect of Yap1 overexpression, mediated by one or more intermediary factors. Yap1 sites were found only four times in the corresponding region of an arbitrary set of 30 genes that were not differentially regulated by Yap1.

Use of a DNA microarray to characterize the transcriptional consequences of mutations affecting the activity of regulatory molecules provides a simple and powerful approach to dissection and characterization of regulatory pathways and net-


works. This strategy also has an important practical application in drug screening. Mutations in specific genes encoding candidate drug targets can serve as surrogates for the ideal chemical inhibitor or modulator of their activity. DNA microarrays can be used to define the resulting signature pattern of alterations in gene expression, and then subsequently used in an assay to screen for compounds that reproduce the desired signature pattern.

DNA microarrays provide a simple and economical way to explore gene expression patterns on a genomic scale. The hurdles to extending this approach to any other organism are minor. The equipment

required for fabricating and using DNA microarrays (9) consists of components that were chosen for their modest cost and simplicity. It was feasible for a small group to accomplish the amplification of more than 6000 genes in about 4 months and, once the amplified gene sequences were in hand, only 2 days were required to print a set of 110 microarrays of 6400 elements each. Probe preparation, hybridization, and fluorescent imaging are also simple procedures. Even conceptually simple experiments, as we described here, can yield vast amounts of information. The value of the information from each experiment of this kind will progressively increase as more is learned about the functions of each gene and as additional experiments define the global changes in gene expression in diverse other natural processes and genetic perturbations. Perhaps the greatest challenge now is to develop efficient methods for organizing, distributing, interpreting, and extracting insights from the large volumes of data these experiments will provide.

## REFERENCES AND NOTES

1. M. Schena, D. Shalon, R. W. Davis, P. O. Brown, *Science* 270, 467 (1995).
2. D. Shalon, S. J. Smith, P. O. Brown, *Genome Res.* 6, 639 (1996).
3. D. Lashkari, *Proc. Natl. Acad. Sci. U.S.A.*, in press.
4. J. DeRisi et al., *Nature Genet.* 14, 457 (1996).
5. D. J. Lockhart et al., *Nature Biotechnol.* 14, 1675 (1996).
6. M. Chee et al., *Science* 274, 610 (1996).
7. M. Johnston and M. Carlson, in *The Molecular Biology of the Yeast Saccharomyces: Gene Expression*, E. W. Jones, J. R. Pringle, J. R. Broach, Eds. (Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1992), p. 193.
8. Primers for each known or predicted protein coding sequence were supplied by Research Genetics. PCR was performed with the protocol supplied by Research Genetics, using genomic DNA from yeast strain S288C as a template. Each PCR product was verified by agarose gel electrophoresis and was deemed correct if the lane contained a single band of appropriate mobility. Failures were marked as such in the database. The overall success rate for a single-pass amplification of 6116 ORFs was ~94.5%.
9. Glass slides (Gold Seal) were cleaned for 2 hours in a solution of 2 N NaOH and 70% ethanol. After rinsing in distilled water, the slides were then treated with a 1:5 dilution of poly-L-lysine adhesive solution (Sigma) for 1 hour, and then dried for 5 min at 40°C in a vacuum oven. DNA samples from 100- $\mu$ l PCR reactions were purified by ethanol purification in 96-well microtiter plates. The resulting precipitates were resuspended in 3 $\times$  standard saline citrate (SSC) and transferred to new plates for arraying. A custom-built arraying robot was used to print on a batch of 110 slides. Details of the design of the microarrayer are available at [cmgm.stanford.edu/pbrown](http://cmgm.stanford.edu/pbrown). After printing, the microarrays were rehydrated for 30 s in a humid chamber and then snap-dried for 2 s on a hot plate (100°C). The DNA was then ultraviolet (UV)-crosslinked to the surface by subjecting the slides to 60 mJ of energy (Stratagene Stratalinker). The rest of the poly-L-lysine surface was blocked by a 15-min incubation in a solution of 70 mM succinic anhydride dissolved in a solution consisting of 315 ml of 1-methyl-2-pyrrolidinone (Aldrich) and 35 ml of 1 M boric acid (pH 8.0). Directly after the blocking reac-



**Fig. 5.** Distinct temporal patterns of induction or repression help to group genes that share regulatory properties. (A) Temporal profile of the cell density, as measured by OD at 600 nm and glucose concentration in the media. (B) Seven genes exhibited a strong induction (greater than ninefold) only at the last timepoint (20.5 hours). With the exception of *IDP2*, each of these genes has a CSRE UAS. There were no additional genes observed to match this profile. (C) Seven members of a class of genes marked by early induction with a peak in mRNA levels at 18.5 hours. Each of these genes contains STRE motif repeats in their upstream promoter regions. (D) Cytochrome c oxidase and ubiquinol cytochrome c reductase genes. Marked by an induction coincident with the diauxic shift, each of these genes contains a consensus binding motif for the HAP2,3,4 protein complex. At least 17 genes shared a similar expression profile. (E) *SAM1*, *GPP1*, and several genes of unknown function are repressed before the diauxic shift, and continue to be repressed upon entry into stationary phase. (F) Ribosomal protein genes comprise a large class of genes that are repressed upon depletion of glucose. Each of the genes profiled here contains one or more RAP1-binding motifs upstream of its promoter. RAP1 is a transcriptional regulator of most ribosomal proteins.

- tion, the bound DNA was denatured by a 2-min incubation in distilled water at ~95°C. The slides were then transferred into a bath of 100% ethanol at room temperature, rinsed, and then spun dry in a clinical centrifuge. Slides were stored in a closed box at room temperature until used.
10. YPD medium (8 liters), in a 10-liter fermentation vessel, was inoculated with 2 ml of a fresh overnight culture of yeast strain DBY7286 (MATa, ura3, GAL2). The fermentor was maintained at 30°C with constant agitation and aeration. The glucose content of the media was measured with a UV test kit (Boehringer Mannheim, catalog number 716251). Cell density was measured by OD at 600-nm wavelength. Aliquots of culture were rapidly withdrawn from the fermentation vessel by peristaltic pump, spun down at room temperature, and then flash frozen with liquid nitrogen. Frozen cells were stored at -80°C.
  11. Cy3-dUTP or Cy5-dUTP (Amersham) was incorporated during reverse transcription of 1.25 µg of polyadenylated [poly(A)<sup>+</sup>] RNA, primed by a dT(16) oligomer. This mixture was heated to 70°C for 10 min, and then transferred to ice. A premixed solution, consisting of 200 U Superscript II (Gibco), buffer, deoxyribonucleoside triphosphates, and fluorescent nucleotides, was added to the RNA. Nucleotides were used at these final concentrations: 500 µM for dATP, dCTP, and dGTP and 200 µM for dTTP. Cy3-dUTP and Cy5-dUTP were used at a final concentration of 100 µM. The reaction was then incubated at 42°C for 2 hours. Unincorporated fluorescent nucleotides were removed by first diluting the reaction mixture with of 470 µl of 10 mM Tris-HCl (pH 8.0)/1 mM EDTA and then subsequently concentrating the mix to ~5 µl, using Centricon-30 microconcentrators (Amicon).
  12. Purified, labeled cDNA was resuspended in 11 µl of 3.5× SSC containing 10 µg poly(dA) and 0.3 µl of 10% SDS. Before hybridization, the solution was boiled for 2 min and then allowed to cool to room temperature. The solution was applied to the microarray under a cover slip, and the slide was placed in a custom hybridization chamber which was subsequently incubated for ~8 to 12 hours in a water bath at 62°C. Before scanning, slides were washed in 2× SSC, 0.2% SDS for 5 min, and then 0.05× SSC for 1 min. Slides were dried before scanning by centrifugation at 500 rpm in a Beckman CS-6R centrifuge.
  13. The complete data set is available on the Internet at [cmgm.stanford.edu/pbrown/explore/index.html](http://cmgm.stanford.edu/pbrown/explore/index.html)
  14. For 95% of all the genes analyzed, the mRNA levels measured in cells harvested at the first and second interval after inoculation differed by a factor of less than 1.5. The correlation coefficient for the comparison between mRNA levels measured for each gene in these two different mRNA samples was 0.98. When duplicate mRNA preparations from the same cell sample were compared in the same way, the correlation coefficient between the expression levels measured for the two samples by comparative hybridization was 0.99.
  15. The numbers and identities of known and putative genes, and their homologies to other genes, were gathered from the following public databases: *Saccharomyces* Genome Database ([genome-www.stanford.edu](http://genome-www.stanford.edu)), Yeast Protein Database ([quest7.proteome.com](http://quest7.proteome.com)), and Munich Information Centre for Protein Sequences ([speedy.mips.biochem.mpg.de/mips/yeast/index.htm](http://speedy.mips.biochem.mpg.de/mips/yeast/index.htm)).
  16. A. Scholer and H. J. Schuller, *Mol. Cell. Biol.* 14, 3613 (1994).
  17. S. Kratzer and H. J. Schuller, *Gene* 161, 75 (1995).
  18. R. J. Haselbeck and H. L. McAlister, *J. Biol. Chem.* 268, 12116 (1993).
  19. M. Fernandez, E. Fernandez, R. Rodicio, *Mol. Gen. Genet.* 242, 727 (1994).
  20. A. Hartig et al., *Nucleic Acids Res.* 20, 5677 (1992).
  21. P. M. Martinez et al., *EMBO J.* 15, 2227 (1996).
  22. J. C. Varela, U. M. Praekelt, P. A. Meacock, R. J. Planta, W. H. Mager, *Mol. Cell. Biol.* 15, 6232 (1995).
  23. H. Ruis and C. Schuller, *Bioessays* 17, 959 (1995).
  24. J. L. Parrou, M. A. Teste, J. Francois, *Microbiology* 143, 1891 (1997).
  25. This expression profile was defined as having an induction of greater than 10-fold at 18.5 hours and less than 11-fold at 20.5 hours.
  26. S. L. Forsburg and L. Guarente, *Genes Dev.* 3, 1166 (1989).
  27. J. T. Olesen and L. Guarente, *ibid.* 4, 1714 (1990).
  28. M. Rosenkrantz, C. S. Kell, E. A. Pennell, L. J. Devenish, *Mol. Microbiol.* 13, 119 (1994).
  29. Single-letter abbreviations for the amino acid residues are as follows: A, Ala; C, Cys; D, Asp; E, Glu; F, Phe; G, Gly; H, His; I, Ile; K, Lys; L, Leu; M, Met; N, Asn; P, Pro; Q, Gln; R, Arg; S, Ser; T, Thr; V, Val; W, Trp; and Y, Tyr. The nucleotide codes are as follows: B-C, G, or T; N-G, A, T, or C; R-A or G; and Y-C or T.
  30. C. Fondrat and A. Kalogeropoulos, *Comput. Appl. Biosci.* 12, 363 (1996).
  31. D. Shore, *Trends Genet.* 10, 408 (1994).
  32. R. J. Planta and H. A. Raue, *ibid.* 4, 64 (1988).
  33. The degenerate consensus sequence VYCYRNNC-MNH was used to search for potential RAP1-binding sites. The exact consensus, as defined by (30), is WACAYCCRTACATYW, with up to three differences allowed.
  34. S. F. Neuman, S. Bhattacharya, J. R. Broach, *Mol. Cell. Biol.* 15, 3187 (1995).
  35. P. Lesage, X. Yang, M. Carlson, *ibid.* 16, 1921 (1996).
  36. For example, we observed large inductions of the genes coding for *PCK1*, *FBP1* [Z. Yin et al., *Mol. Microbiol.* 20, 751 (1996)], the central glyoxylate cycle gene *ICL1* [A. Scholer and H. J. Schuller, *Curr. Genet.* 23, 375 (1993)], and the "aerobic" isoform of acetyl-CoA synthase, *ACS1* [M. A. van den Berg et al., *J. Biol. Chem.* 271, 28953 (1996)], with concomitant down-regulation of the glycolytic-specific genes *PYK1* and *PFK2* [P. A. Moore et al., *Mol. Cell. Biol.* 11, 5330 (1991)]. Other genes not directly involved in carbon metabolism but known to be induced upon nutrient limitation include genes encoding cytosolic catalase *CTT1* [P. H. Bissinger et al., *ibid.* 9, 1309 (1989)] and several genes encoding small heat-shock proteins, such as *HSP12*, *HSP26*, and *HSP42* [I. Farkas et al., *J. Biol. Chem.* 266, 15602 (1991); U. M. Praekelt and P. A. Meacock, *Mol. Gen. Genet.* 223, 97 (1990); D. Wotton et al., *J. Biol. Chem.* 271, 2717 (1996)].
  37. The levels of induction we measured for genes that were expressed at very low levels in the uninduced state (notably, *FBP1* and *PCK1*) were generally lower than those previously reported. This discrepancy was likely due to the conservative background subtraction method we used, which generally resulted in overestimation of very low expression levels (46).
  38. Cross-hybridization of highly related sequences can also occasionally obscure changes in gene expression, an important concern where members of gene families are functionally specialized and differentially regulated. The major alcohol dehydrogenase genes, *ADH1* and *ADH2*, share 88% nucleotide identity. Reciprocal regulation of these genes is an important feature of the diauxic shift, but was not observed in this experiment, presumably because of cross-hybridization of the fluorescent cDNAs representing these two genes. Nevertheless, we were able to detect differential expression of closely related isoforms of other enzymes, such as *HXK1/HXK2* (77% identical) [P. Herrero et al., *Yeast* 11, 137 (1995)], *MLS1/DAL7* (73% identical) (20), and *PGM1/PGM2* (72% identical) [D. Oh, J. E. Hopper, *Mol. Cell. Biol.* 10, 1415 (1990)], in accord with previous studies. Use in the microarray of deliberately selected DNA sequences corresponding to the most divergent segments of homologous genes, in lieu of the complete gene sequences, should relieve this problem in many cases.
  39. F. E. Williams, U. Varanasi, R. J. Trumbly, *Mol. Cell. Biol.* 11, 3307 (1991).
  40. D. Tzamas and K. Struhl, *Nature* 369, 758 (1994).
  41. Differences in mRNA levels between the *tup1Δ* and wild-type strain were measured in two independent experiments. The correlation coefficient between the complete sets of expression ratios measured in these duplicate experiments was 0.83. The concordance between the sets of genes that appeared to be induced was very high between the two experiments. When only the 355 genes that showed at least a twofold increase in mRNA in the *tup1Δ* strain in either of the duplicate experiments were compared, the correlation coefficient was 0.82.
  42. The *tup1Δ* mutation consists of an insertion of the *LEU2* coding sequence, including a stop codon, between the ATG of *TUP1* and an Eco RI site 124 base pairs before the stop codon of the *TUP1* gene.
  43. L. R. Kowalski, K. Kondo, M. Inouye, *Mol. Microbiol.* 15, 341 (1995).
  44. M. Viswanathan, G. Muthukumar, Y. S. Cong, J. Lenard, *Gene* 148, 149 (1994).
  45. D. Hirata, K. Yano, T. Miyakawa, *Mol. Gen. Genet.* 242, 250 (1994).
  46. A. Gutierrez, L. Caramelo, A. Prieto, M. J. Martinez, A. T. Martinez, *Appl. Environ. Microbiol.* 60, 1783 (1994).
  47. A. Muheim et al., *Eur. J. Biochem.* 195, 369 (1991).
  48. J. A. Wemmie, M. S. Szczypka, D. J. Thiele, W. S. Moye-Rowley, *J. Biol. Chem.* 269, 32592 (1994).
  49. Microarrays were scanned using a custom-built scanning laser microscope built by S. Smith with software written by N. Ziv. Details concerning scanner design and construction are available at [cmgm.stanford.edu/pbrown](http://cmgm.stanford.edu/pbrown). Images were scanned at a resolution of 20 µm per pixel. A separate scan, using the appropriate excitation line, was done for each of the two fluorophores used. During the scanning process, the ratio between the signals in the two channels was calculated for several array elements containing total genomic DNA. To normalize the two channels with respect to overall intensity, we then adjusted photomultiplier and laser power settings such that the signal ratio at these elements was as close to 1.0 as possible. The combined images were analyzed with custom-written software. A bounding box, fitted to the size of the DNA spots in each quadrant, was placed over each array element. The average fluorescent intensity was calculated by summing the intensities of each pixel present in a bounding box, and then dividing by the total number of pixels. Local area background was calculated for each array element by determining the average fluorescent intensity for the lower 20% of pixel intensities. Although this method tends to underestimate the background, causing an underestimation of extreme ratios, it produces a very consistent and noise-tolerant approximation. Although the analog-to-digital board used for data collection possesses a wide dynamic range (12 bits), several signals were saturated (greater than the maximum signal intensity allowed) at the chosen settings. Therefore, extreme ratios at bright elements are generally underestimated. A signal was deemed significant if the average intensity after background subtraction was at least 2.5-fold higher than the standard deviation in the background measurements for all elements on the array.
  50. In addition to the 17 genes shown in Table 1, three additional genes were induced by an average of more than threefold in the duplicate experiments, but in one of the two experiments, the induction was less than twofold (range 1.6- to 1.9-fold).
  51. We thank H. Bennett, P. Spellman, J. Ravetto, M. Eisen, R. Pillai, B. Dunn, T. Ferea, and other members of the Brown lab for their assistance and helpful advice. We also thank S. Friend, D. Botstein, S. Smith, J. Hudson, and D. Dolginow for advice, support, and encouragement; K. Struhl and S. Chatterjee for the *Tup1* deletion strain; L. Fernandes for helpful advice on *Yap1*; and S. Klapholz and the reviewers for many helpful comments on the manuscript. Supported by a grant from the National Human Genome Research Institute (NHGRI) (HG00450), and by the Howard Hughes Medical Institute (HHMI). J.D.R. was supported by the HHMI and the NHGRI. V.R. was supported in part by an Institutional Training Grant in Genome Science (T32 HG00044) from the NHGRI. P.O.B. is an associate investigator of the HHMI.

5 September 1997; accepted 22 September 1997



# The New York Times

ON THE WEB

October 2, 2003, Thursday

BUSINESS/FINANCIAL DESK

## Human Genome Placed on Chip; Biotech Rivals Put It Up for Sale

By ANDREW POLLACK (NYT) 1030 words

The genome on a chip has arrived.

Melding high technology with biology, several companies are rushing to sell slivers of glass or nylon, some as small as postage stamps, packed with pieces of all 30,000 or so known human genes.

The new products will allow scientists to scan all genes in a human tissue sample at once, to determine which genes are active, a job that previously required two or more chips. The whole-genome chips will lower the cost and increase the speed of a widely used test that has transformed biomedical research in the last few years.

"It's sort of a milestone event, very similar to generating an integrated circuit of the genome," said Stephen P. A. Fodor, the chief executive of Affymetrix Inc., the leading seller of gene chips, which are also called microarrays.

Affymetrix, based in Santa Clara, Calif., is expected to announce today that it is accepting orders for its whole-genome chip.

The announcement seems timed to steal some thunder from the rival Agilent Technologies, which is based in nearby Palo Alto. Agilent is to be the host of an analyst meeting today and it plans to announce then that it has started shipping test versions of its whole-genome chip.

Applied Biosystems of Foster City, Calif., a unit of the Applera Corporation, started the race in July with an announcement that it would have a whole-genome chip out by the end of this year. NimbleGen Systems, a small company in Madison, Wis., announced a few days later that it had a genome on a chip that it was not selling but that it was using to run tests for customers.

Gene chips, which detect genes that are active, meaning they are being used to make a protein, have become essential tools. Scientists try to understand the genetic mechanisms of disease by seeing which genes are turned on in, say, a sick kidney or lung compared with those active in a healthy organ. Pharmaceutical companies look at gene activity patterns to try to predict the effects of drugs.

Scientists have found that tumors that look the same under the microscope can differ in terms of which genes are active. So studying gene patterns could become a way to discriminate between deadly and not-so-deadly tumors, or to predict which drug will work best for a particular patient.

Still, even some vendors conceded that the change from two chips to one is more symbolic than revolutionary.

"You can do just as good science with two chips, it costs you a little more," said Roland Green, the vice president for research and development at NimbleGen.

Some scientists questioned whether the chips really have all human genes, because the exact number and identities of all the genes is not known.

The advent of the genome on a chip is, however, evidence that biotechnology, to the extent that it uses electronics, is experiencing some of the rapid progress that has made semiconductors and computers continuously cheaper and smaller.

"One of the effects everyone is looking for in the genomics area is Moore's law -- more data, less money," said Doug Dolginow, an executive vice president at Gene Logic, which sells data from gene chip studies to pharmaceutical companies. "This is a step in that direction."

Moore's law states that the number of transistors on a semiconductor chip doubles every 18 months.

Affymetrix's gene chips are, in fact, made with the same techniques used to make semiconductor chips. In the mid-1990's, the company came out with a set of five chips covering what was then known of the human genome. After the human genome sequence was virtually completed in 2000, the company developed a two-chip set with all the known genes. Now it has the single chip, which some scientists say will be more convenient.

"We like to be able to look at all genes at one time to get a global view of what's going on," said John R. Walker, who runs gene chip operations at the Genomics Institute of the Novartis Research Foundation in San Diego.

Costs should also be lower. Gene chips have been so expensive that many academic scientists still make their own rather than buy them. Affymetrix said it would sell its whole-genome chips for \$300 to \$500 each, depending on volume, little more than half the price of the two-chip set. The other companies have not announced prices.

For Affymetrix, a successful whole-genome chip "is essential for them to maintain their dominance" of high-end microarrays, said Edward A. Tenthoff, an analyst at U.S. Bancorp Piper Jaffray. Affymetrix had total product sales in 2002 of about \$250 million, and a company spokesman said that human genome chips are its top-selling product.

Mr. Tenthoff, who recommends Affymetrix stock, said the company's sales growth rate had moderated as it faces tougher competition. Agilent, a spinoff of Hewlett-Packard that makes its gene chips by printing DNA components onto glass slides using ink jet printers, has gained share, he said. Applied Biosystems, the largest maker of genomics equipment over all, will be

entering the microarray segment of the business with its whole-genome chip, emphasizing the connection of that product to the others it offers, including the gene database developed by its sister company, Celera Genomics.

Jeffrey Trent, scientific director of the Translational Genomics Research Institute in Phoenix, said that while whole-genome chips are useful for medical discovery, the biggest growth of the market will be for chips that can be used by doctors to do diagnoses. And whole-genome chips are too cumbersome for that, he said. Rather, once scientists use the whole-genome chips to find particular genes that are associated with, say, tumor aggressiveness or drug effectiveness, he said, they will then make smaller and cheaper chips containing just those genes for use in diagnosis.



Agilent | Agilent Technologies ships whole human genome on single microarray to gene e...



About Agilent | Products & Services | Industries | International | Online Stores

[Worldwide Home](#) > [About Agilent](#) > [News@Agilent](#) > [Press Releases](#)

News@Agilent

## Agilent Technologies ships whole human genome on single microarray to gene expression customers for evaluation

Company to introduce first commercial whole human microarray by end of year

PALO ALTO, Calif., Oct. 2, 2003

### Press Releases

- ▶ [Communi](#)
- ▶ [Corporate](#)
- ▶ [Electronic](#)
- ▶ [Life Scien](#)  
[Chemical](#)

### ▶ [Archives](#)

### [Search Agile](#)

### [Quick Links](#)

[Jump to pa](#)

Agilent Technologies Inc. (NYSE: A) today announced it has shipped whole human-genome microarrays to customers for testing and evaluation. The whole genome microarray is based on Agilent's new double-density format, which can accommodate 44,000 features on a single 1" x 3" glass-slide microarray. The new platform enables drug-discovery and disease researchers to perform whole-genome screening at a lower cost and with higher reproducibility.

"This is an important step toward our release of the first whole human-genome microarray product, which is expected to be available for order before the end of the year," said Barney Saunders, vice president and general manager of Agilent's BioResearch Solutions Unit. "Customers have long wanted a one-sample, one-chip format with the increased sensitivity associated with 60-mer probes. The cost savings and high-quality performance make this product a compelling alternative for scientists who make their own microarrays."

Agilent's microarrays are based on the industry-standard 1" x 3" (25mm x 75mm) format, which is compatible with most commercial microarray scanners. All Agilent commercial microarrays are developed using content from public databases and proprietary sources, with full sequence and annotation information made available to customers. Gene sequences for probes are developed using algorithms and then validated empirically through iterative wet-lab testing procedures. The result is a microarray comprised of functionally validated probes, with the most up-to-date and comprehensive genome information commercially available.

Advantages of the double-density format include:

- Lower cost. Not only is one microarray less expensive than two, it requires fewer reagents and reduces instrumentation demands.
- Streamlined workflow. Researchers need prepare and process only one microarray instead of two. This also results in fewer steps in the subsequent data analysis.
- Greater reproducibility. Use of a single microarray further reduces unnecessary variability in experimental conditions.
- Smaller sample use. A smaller quantity of sample material is required to perform an experiment.

### Availability

Agilent's Whole Human Genome Microarray is expected to be available for order by the end of the year.

### About Agilent Technologies

Agilent Technologies Inc. (NYSE: A) is a global technology leader in communications, electronics, life sciences and chemical analysis. The company's 30,000 employees serve customers in more than 110 countries. Agilent had net revenue of \$6 billion in fiscal year 2002. Information about Agilent is available

on the Web at [www.agilent.com](http://www.agilent.com).

#### Forward-Looking Statements

This news release contains forward-looking statements (including, without limitation, statements relating to Agilent's expectation that its whole-genome microarray platform will be available for order before the end of 2003) that involve risks and uncertainties that could cause results to differ materially from management's current expectations. These and other risks are detailed in the company's filings with the Securities and Exchange Commission, including its Annual Report on Form 10-K for the year ended Oct. 31, 2002, its Quarterly Report on Form 10-Q for the quarter ended July 31, 2003 and its Current Report on Form 8-K filed Aug. 18, 2003. The company assumes no obligation to update the information in this press release.

###

#### Contact:

Christina Maehr  
+1 408 553 7205  
[christina\\_maehr@agilent.com](mailto:christina_maehr@agilent.com)

To send feedback about this site: [Contact Webmaster](#)

© Agilent 2000-2003

[Terms of Use](#)

[Privacy](#)

Today's News

## Today's News

### Affymetrix Announces Commercial Launch of Single Array for Human Genome Expression Analysis



#### AFFYMETRIX GENECHIP(R) BRAND HUMAN GENOME U133 PLUS 2.0 ARRAY

Affymetrix GeneChip(R) Brand Human Genome U133 Plus 2.0 Array.  
(PRNewsFoto)[AS]  
SANTA CLARA, CA USA 10/02/2003

Website

More Than 1 Million Probes Analyze Expression Levels of Nearly 50,000 RNA Transcripts and Variants on a Single Array the Size of a Thumbnail

SANTA CLARA, Calif., Oct. 2 /PRNewswire/ -- Affymetrix, Inc., (Nasdaq: AFFX) announced today that it is taking orders for its new GeneChip(R) brand Human Genome U133 Plus 2.0 Array, offering researchers the protein-coding content of the human genome on a single commercially available catalog microarray. The HG-U133 Plus 2.0 Array analyzes the expression level of nearly 50,000 RNA transcripts and variants with 22 different probes per transcript, providing superior data quality unmatched by technologies using a single probe per transcript.

(Photo: <http://www.newscom.com/cgi-bin/prnh/20031002/SFTH021> )

"With about 1.3 million probes on a chip the size of a human thumbnail, the Human Plus Array represents a leap in array technology data capacity, and further demonstrates the unique power and potential of our technology to explore vast areas of the genome," said Trevor J. Nicholls, Ph.D., Chief Commercial Officer. "Multiple independent measurements for each transcript ensure that our data quality remains the industry standard, even as our data capacity increases dramatically."

The HG-U133 Plus 2.0 Array, which will ship in October, combines the content of the previous HG-U133 two-array set with nearly 10,000 new probe sets representing about 6,500 new genes, for a total of nearly 50,000 RNA transcripts and variants. This new information, verified against the latest version of the publicly available genome map, provides researchers the most comprehensive and up-to-date genome-wide gene expression analysis. The probe design strategy of the HG-U133 Plus 2.0 Array is identical to the previous HG-U133 Set, providing very strong data concordance between the two products. With more than double the data capacity of the previous-generation Affymetrix human product, the HG-U133 Plus 2.0 Array can significantly cut processing and analysis time for scientists in the lab, freeing up valuable resources and accelerating research.

The HG-U133 Plus 2.0 Array sets a new standard for the number of genes and transcripts on any commercially available single array for human gene



expression analysis, while maintaining Affymetrix' unrivaled data quality. The HG-U133 Plus 2.0 Array uses 22 independent measures to detect the hybridization of each transcript on the array, 1.3 million data points in all, more than 30 times that of any other microarray technology. Using multiple, independent measurements provides optimal sensitivity and specificity, and the most accurate, consistent and statistically significant results possible.

"More data points produce more reliable results and ultimately, enable better science," said Nicholls. "Our powerful probe set strategy gives our customers the assurance that their array results actually reflect what's in their sample."

Affymetrix is also launching an updated 11-micron version of its popular 18-micron HG-U133A Array called the GeneChip HG-U133A 2.0 Array. The reduced feature size on this new design means researchers can use smaller sample volumes than on the previous 18-micron array without compromising performance. This new array represents over 20,000 transcripts that can be used to explore human biology and disease processes. All probe sets represented on the original GeneChip HG-U133A Array are identically replicated on the GeneChip HG-U133A 2.0 Array.

More information on the design of the HG-U133 Plus 2.0 Array and the HG-U133A 2.0 Array may be found on the Affymetrix website at <http://www.affymetrix.com>.

Affymetrix will be presenting further information on this and other products at the BioTechnica trade show in Hanover, Germany on Oct. 7-9, 2003. The Company will also hold a press conference on Oct. 7, from 11 a.m. to 12 p.m. at the show regarding the new Human Genome U133 Plus 2.0 Array. If you would like to attend this press conference, please contact Caroline Stupnicka at [c.stupnicka@northbankcommunications.com](mailto:c.stupnicka@northbankcommunications.com).

#### About Affymetrix:

Affymetrix is a pioneer in creating breakthrough tools that are driving the genomic revolution. By applying the principles of semiconductor technology to the life sciences, Affymetrix develops and commercializes systems that enable scientists to improve the quality of life. The Company's customers include pharmaceutical, biotechnology, agrichemical, diagnostics and consumer products companies as well as academic, government and other non-profit research institutes. Affymetrix offers an expanding portfolio of integrated products and services, including its integrated GeneChip platform, to address growing markets focused on understanding the relationship between genes and human health. Additional information on Affymetrix can be found at <http://www.affymetrix.com>.

All statements in this press release that are not historical are "forward-looking statements" within the meaning of Section 21E of the Securities Exchange Act as amended, including statements regarding Affymetrix' "expectations," "beliefs," "hopes," "intentions," "strategies" or the like. Such statements are subject to risks and uncertainties that could cause actual results to differ materially for Affymetrix from those projected, including, but not limited to risks of the Company's ability to achieve and sustain higher levels of revenue, higher gross margins, reduced operating expenses, uncertainties relating to technological approaches, manufacturing, product development, market acceptance (including uncertainties relating to product development and market acceptance of the GeneChip HG-U133 Human Plus 2.0 Array and the HG-U133A 2.0), personnel retention, uncertainties related to cost and pricing of Affymetrix products, dependence on collaborative partners, uncertainties relating to sole source suppliers, uncertainties relating to FDA and other regulatory approvals, competition, risks relating to intellectual property of others and the uncertainties of patent protection and litigation. These and other risk factors are discussed in Affymetrix' Form 10-K for the

year ended December 31, 2002 and other SEC reports, including its Quarterly Reports on Form 10-Q for subsequent quarterly periods. Affymetrix expressly disclaims any obligation or undertaking to release publicly any updates or revisions to any forward-looking statements contained herein to reflect any change in Affymetrix' expectations with regard thereto or any change in events, conditions, or circumstances on which any such statements are based.

NOTE: Affymetrix, the Affymetrix logo, and GeneChip and are registered trademarks owned or used by Affymetrix, Inc.

**SOURCE Affymetrix, Inc.**

**Web Site: <http://www.affymetrix.com>**

**Photo Notes: NewsCom:**

**<http://www.newscom.com/cgi-bin/prnh/20031002/SFTH021> AP Archive:**

**<http://photoarchive.ap.org> PRN Photo Desk,**

**[photodesk@prnewswire.com](mailto:photodesk@prnewswire.com)**

---

**Issuers of news releases and not PR Newswire are solely responsible for the accuracy of the content.**

**More news from PR Newswire...**

**Copyright © 1996-2002 PR Newswire Association LLC. All Rights Reserved.  
A United Business Media company.**

# Macroresults through Microarrays

John C. Rockett, Reproductive Toxicology Division (MD-72), National Health and Environmental Effects Research Laboratory, Office of Research and Development, US Environmental Protection Agency, Research Triangle Park, 2525 East Highway 54, Durham, NC 27711, USA; tel: +1 919 541 2071, fax: +1 919 541 4017, e-mail: rockett.john@epa.gov

The third enactment of Cambridge Healthtech Institute's *Macroresults through Microarrays* meeting was held in Boston (MA, USA) from 29 April–1 May 2002. The subtheme of this year's meeting was 'advancing drug discovery', a widely touted application for array technology.

## The evolution of microarrays

If you were asked 'Who first conceived of the idea of microarrays', who would come to mind? Mark Schena perhaps, first author of the seminal 1995 paper on cDNA arrays [1]? Maybe Pat Brown, Schena's then supervisor? Or perhaps Stephen Fodor, the primary driver behind Affymetrix's (<http://www.affymetrix.com>) oligonucleotide-based platform [2]. Brits might even chant the name of Ed Southern [3]. Well, according to Roger Ekins (University College London Medical School; <http://www.ucl.ac.uk/medicine/>) all these answers would be wrong. It was in fact Ekins and his colleagues who first conceived of and patented 'a new generation of ultrasensitive, miniaturized assays for protein and DNA–RNA measurement based on the use of microarrays' in the mid 1980s [4]. The concept and potential of array technology was more fully described in a later publication, in which Ekins *et al.* [5] concluded that antibody microspots of  $\sim 50 \mu\text{m}^2$  could be achieved, and that as many as 2 million different immunoassays could, in principle, be accommodated on a surface area of  $1 \text{ cm}^2$ .

## Technological innovation

In practice, it took a different biological molecule (DNA), a different research

group, and a leap into microfabrication technology to even begin approaching these kinds of densities [Affymetrix patent 6045996 talks of one million spots  $\text{cm}^{-2}$ ]. Of course, advancing technology is one of the driving engines behind the genomics juggernaut, and we are already seeing '4th generation' machines for fabricating DNA chips. If the company representatives at this meeting are to be believed (and their cases seemed strong), spotting is out, and *in situ* fabrication of oligonucleotide-based 'iterative custom arrays' is in. Whether you go with the Combimatrix's (<http://www.combimatrix.com>) electrochemically directed synthesis and detection system, febit's (<http://www.febit.com>) Geniom® technology, or Nimblegen's (<http://www.nimblegen.com>) Maskless Array Synthesizer technology is a matter of personal choice. However, each of these machines provides the flexibility to design variable length oligonucleotide probes from sequences inputted by the user, and then perform *in situ* synthesis of an array. Each system also boasts unique advantages. For example, Combimatrix's biological array processor is a semiconductor coated with a 3D layer of porous material in which DNA, RNA, peptides or small molecules can be synthesized or immobilized within discrete test sites, while febit's Geniom One® is a fully integrated gene-expression analysis system with minimal user hands-on time – the probe sequences are programmed, the RNA samples inserted, and the gene expression data is pumped out a few hours later.

## Cell- and tissue-based arrays

Array technology is in most people's minds firmly linked with gene-expression profiling. Fewer are aware that cell- and tissue-based arrays have been developed, and how they can provide a vital extra dimension to research. In support of this, Barry Bochner gave an update on the cell-based array system that Biolog (<http://www.biolog.com>) has produced for simultaneously measuring the effects of one gene in the cell under thousands of growth conditions (see [6] for further details). David Walt (Tufts University; <http://www.tufts.edu/>) is developing single live cell arrays using optical imaging fiber (OIF) technology. An array of microwells is fabricated on the face of an OIF at densities of up to 10 million wells  $\text{cm}^{-2}$ . Cells are then added to the wells and disperse at an average of one cell per well. Physiological and genetic responses of each cell are measured via fluorescence produced by reporter genes (e.g. *lacZ*, *gfp*). Assays performed so far include yeast live or dead cell assay, microenvironment pH and  $\text{O}_2$  measurements, promotor responses using the *lacZ* and *phoA* reporter genes, and protein–protein interactions using the yeast two-hybrid system. The main advantage of this system is that the cells remain alive during the assay, which means a real-time timecourse can be performed and/or the array passed from sample to sample. This would be useful in, for example, the scanning of a combinatorial drug library for specific physiological effects.

Tissue arrays are a useful complementary technology to DNA arrays because they can be used to help validate and

understand the biological and medical significance of gene changes discovered using standard DNA arrays. For example, an array of tumor tissues can be screened for the protein (using immunohistochemistry), message (using *in situ* hybridization) and copy number (using comparative genomic hybridization) of a gene of interest, to determine if expression of the gene (or lack thereof) is related in any way to survival. They can also be used to predict the probability of clinical failure of lead compounds as a result of toxicity by evaluating the distribution of the drug targets in normal tissue. Spyro Mousses and his co-workers at the National Human Genome Research Institute (<http://www.nhgri.nih.gov/index.html>) have built such arrays, including a multi-tumor array (~5000 specimens, and sections from 36 normal and 800 metastatic tissues) and a normal tissue array (76 tissue and 332 cell types).

#### The problem with proteins

It has been said that genomics tells us what might happen, transcriptomics indicates what should happen, and proteomics shows what is happening. The impact of functional proteomics on pharmaceutical R&D is rapidly increasing, and protein arrays are being used increasingly in both basic and applied research. Their use lies not only in comparative protein expression and interaction profiling, but also in diagnostics and drug discovery. However, an increasing number of researchers have found that protein arrays, like their cousins the DNA arrays, present several practical obstacles relating to their production and use. For example, in using *Escherichia coli* to produce recombinant eukaryotic proteins from a single expression vector, multiple protein products are often produced, suggesting mixes of truncated or otherwise altered proteins. There is also the obvious concern that the proteins might not be modified in a similar manner to

eukaryotic systems. Also, an optimal method for depositing and binding proteins to the selected substrate is yet to be determined, as is the best way to ensure that they are bound in a correctly folded, active conformation.

Several companies have been addressing these problems. Prolinx (<http://www.prolinxinc.com>) is one such company, and Karin Hughes described their Versalinx™ chemistry for producing protein, peptide and small-molecule arrays. Versalinx™ uses solution-phase conjugation followed by immobilization, resulting in functional orientation of proteins and peptides on the substrate surface. It also offers the valuable additional benefit of exhibiting low non-specific binding. Sense Proteomic (<http://www.senseproteomic.com>) is also among those addressing these problems to develop robust protein arrays for drug discovery and clinical applications and has developed functional protein array formats based on specific disease tissues. Subtractive hybridization is used to identify genes with altered expression in breast tumor and cystic fibrosis compared to normal tissue. A high throughput cloning strategy (COVET™) is then used to produce libraries of genes that are tagged, cloned, expressed, purified and finally immobilized on glass slides. Initial validation studies have shown that the vast majority of the immobilized proteins do indeed retain biological function.

Stefan Schmidt and his company (GPC Biotech; <http://www.gpcbiotech.de>) have moved past the platform development stage and, with their focus firmly on drug discovery, are currently developing kinase-profiling arrays. Kinases are important targets for pharmaceutical drug discovery and therapy, and GPC's aim is to simultaneously detect multiple kinases, obtain activity profiles for different cell types, or analyze the ability of drug candidates to inhibit kinase activity. To do this, recombinant kinase substrates are immobilized on

membranes, incubated with purified kinase, and the substrates measured for the degree of phosphorylation.

#### Summary

Meetings like this, packed with exciting discoveries and intriguing and interesting innovation, heavily emphasize the pace at which biotechnology is advancing, to the extent that the number of options for genomic and proteomic researchers can become overwhelming. Although data analysis is perhaps the greatest current concern for array users, an increasing challenge will be to determine the approaches and technology that really work, and to do it in a timely manner.

#### References

- 1 Schena, M. *et al.* (1995) Quantitative monitoring of gene expression patterns with a complementary DNA microarray. *Science* 270, 467-470
- 2 Fodor, S.P. *et al.* (1991) Light-directed, spatially addressable parallel chemical synthesis. *Science* 251, 767-773
- 3 Southern, E.M. *et al.* (1992) Analyzing and comparing nucleic acid sequences by hybridization to arrays of oligonucleotides: evaluation using experimental models. *Genomics* 13, 1008-1017
- 4 Ekins, R.P. (1987) US Patent Application 8 803 000
- 5 Ekins, R. *et al.* (1989) High specific activity chemiluminescent and fluorescent markers: their potential application to high sensitivity and 'multi-analyte' immunoassays. *J. Biolum. Chemilum.* 4, 59-78
- 6 Rockett, J.C. (2002) Chip, chip, array! Three chips for post-genomic research. *Drug Discov. Today* 7, 458-459

#### Acknowledgements

I would like to thank Mary Ann Brown (Cambridge Healthtech Institute) and David Dix (US EPA) for critical review of this manuscript prior to submission. This document has been reviewed in accordance with US Environmental Protection Agency policy and approved for publication. Mention of companies, trade names or products does not signify endorsement of such by the EPA.